

79

19



Europäisches Patentamt

Eur pean Patent Office

Offic uropéen des br vets

11 Publication number:

0 130 756  
A1

12

# EUROPEAN PATENT APPLICATION

21 Application number: 84304252.4

22 Date of filing: 22.06.84

51 Int. Cl.: C 12 N 15/00, C 12 N 9/14,  
C 12 N 9/54, C 11 D 3/386,  
C 11 D 7/42  
// C12R1/125

30 Priority: 24.06.83 US 507419  
29.05.84 US 614612  
29.05.84 US 614615  
29.05.84 US 614616  
29.05.84 US 614617  
29.05.84 US 614491

43 Date of publication of application: 09.01.85  
Bulletin 85/2

84 Designated Contracting States: AT BE CH DE FR GB IT  
LI LU NL SE

71 Applicant: GENENTECH, INC., 460 Point San Bruno  
Boulevard, South San Francisco California 94080 (US)

72 Inventor: Bott, Richard Ray, 3032 Hillside Drive,  
Burlingame California 94010 (US)  
Inventor: Ferrari, Eugenio, 75 Morningside Drive, Daly  
City California 94015 (US)  
Inventor: Wells, James Allen, 65 Otay Avenue, San  
Mateo California 94403 (US)  
Inventor: Estell, David Aaron, 250 Diablo Avenue,  
Mountain View California 94043 (US)  
Inventor: Henner, Dennis James, 297 Angellita Avenue,  
Pacifica California 94404 (US)

74 Representative: Armitage, Ian Michael et al, MEWBURN  
ELLIS & CO. 2/3 Cursitor Street, London EC4A 1BQ (GB)

54 Prokaryotic carbonyl hydrolases, methods, DNA, vectors and transformed hosts for producing them, and detergent  
compositions containing them.

57 Methods and vectors are provided for the production of  
prokaryotic carbonyl hydrolases in recombinant systems. DNA  
which encodes such hydrolases are mutated at predetermined  
regions by known methods or by a novel point mutagenesis  
method in order to generate mutant hydrolases. Particular  
point mutations in carbonyl hydrolases such as subtilisin result  
in modifications of oxidation stability, Km, Kcat, Kcat/Km ratio  
substrate specificity, specific activity or pH-activity profiles.  
These mutated hydrolases are particularly useful in laundry  
compositions. Mutations in the genes encoding the subtilisin or  
neutral protease of bacillus yield substantially normally sporu-  
lating bacillus strains which are incapable of excreting subtil-  
isin or neutral protease. Such strains are useful in the recombi-  
nant synthesis of heterologous proteins.

EP 0 130 756 A1

0130756

100/157,241,242,243,247,248

5

10

PROCARYOTIC CARBONYL HYDROLASES, METHODS, DNA,  
VECTORS AND TRANSFORMED HOSTS FOR PRODUCING  
THEM, AND DETERGENT COMPOSITIONS CONTAINING THEM

15

20 Background

This invention relates to the production and manipulation of proteins using recombinant techniques in suitable hosts. More specifically, the invention relates to the production of procaryotic proteases such as subtilisin and neutral protease using recombinant microbial host cells, to the synthesis of heterologous proteins by microbial hosts, and to the directed mutagenesis of enzymes in order to modify the characteristics thereof.

30

Various bacteria are known to secrete proteases at some stage in their life cycles. Bacillus species produce two major extracellular proteases, a neutral protease (a metalloprotease inhibited by EDTA) and an alkaline protease (or subtilisin, a serine endoprotease).

35

Both generally are produced in greatest quantity after the exponential growth phase, when the culture enters stationary phase

and begins the process of sporulation. The physiological role of these two proteases is not clear. They have been postulated to play a role in sporulation (J. Hoch, 1976, "Adv. Genet." 18:69-98; P. Piggot et al., 1976, "Bact. Rev." 40:908-962; and F. Priest, 5 1977, "Bact. Rev." 41:711-753), to be involved in the regulation of cell wall turnover (L. Jolliffe et al., 1980, "J. Bact." 141:1199-1208), and to be scavenger enzymes (Priest, Id.). The regulation of expression of the protease genes is complex. They appear to be coordinately regulated in concert with sporulation, 10 since mutants blocked in the early stages of sporulation exhibit reduced levels of both the alkaline and neutral protease. Additionally, a number of pleiotropic mutations exist which affect the level of expression of proteases and other secreted gene products, such as amylase and levansucrase (Priest, Id.).

15

Subtilisin has found considerable utility in industrial and commercial applications (see U.S. Patent No. 3,623,957 and J. Millet, 1970, "J. Appl. Bact." 33:207). For example, subtilisins and other proteases are commonly used in detergents to enable 20 removal of protein-based stains. They also are used in food processing to accommodate the proteinaceous substances present in the food preparations to their desired impact on the composition.

Classical mutagenesis of bacteria with agents such as radiation 25 or chemicals has produced a plethora of mutant strains exhibiting different properties with respect to the growth phase at which protease excretion occurs as well as the timing and activity levels of excreted protease. These strains, however, do not approach the ultimate potential of the organisms because the mutagenic process is 30 essentially random, with tedious selection and screening required to identify organisms which even approach the desired characteristics. Further, these mutants are capable of reversion to the parent or wild-type strain. In such event the desirable property is lost. The probability of reversion is unknown when dealing with random 35 mutagenesis since the type and site of mutation is unknown or poorly

0992Y

characterized. This introduces considerable uncertainty into the industrial process which is based on the enzyme-synthesizing bacterium. Finally, classical mutagenesis frequently couples a desirable phenotype, e.g., low protease levels, with an undesirable character such as excessive premature cell lysis.

Special problems exist with respect to the proteases which are excreted by *Bacillus*. For one thing, since at least two such proteases exist, screening for the loss of only one is difficult. Additionally, the large number of pleiotropic mutations affecting both sporulation and protease production make the isolation of true protease mutations difficult.

Temperature sensitive mutants of the neutral protease gene have been obtained by conventional mutagenic techniques, and were used to map the position of the regulatory and structural gene in the *Bacillus subtilis* chromosome (H. Uehara et al., 1979, "J. Bact." 139:583-590). Additionally, a presumed nonsense mutation of the alkaline protease gene has been reported (C. Roitsch et al., 1983, "J. Bact." 155:145-152).

*Bacillus* temperature sensitive mutants have been isolated that produce inactive serine protease or greatly reduced levels of serine protease. These mutants, however, are asporogenous and show a reversion frequency to the wild-type of about from  $10^{-7}$  to  $10^{-8}$  (F. Priest, Id. p. 719). These mutants are unsatisfactory for the recombinant production of heterologous proteins because asporogenous mutants tend to lyse during earlier stages of their growth cycle in minimal medium than do sporogenic mutants, thereby prematurely releasing cellular contents (including intracellular proteases) into the culture supernatant. The possibility of reversion also is undesirable since wild-type revertants will contaminate the culture supernatant with excreted proteases.

Bacillus sp. have been proposed for the expression of heterologous proteins, but the presence of excreted proteases and the potential resulting hydrolysis of the desired product has retarded the commercial acceptance of Bacillus as a host for the expression of heterologous proteins. Bacillus megaterium mutants have been disclosed that are capable of sporulation and which do not express a sporulation-associated protease during growth phases. However, the assay employed did not exclude the presence of other proteases, and the protease in question is expressed during the sporulation phase (C. Loshon et al., 1982, "J. Bact." 150:303-311). This, of course, is the point at which heterologous protein would have accumulated in the culture and be vulnerable. It is an objective herein to construct a Bacillus strain that is substantially free of extracellular neutral and alkaline protease during all phases of its growth cycle and which exhibits substantially normal sporulation characteristics. A need exists for non-revertible, otherwise normal protease deficient organisms that can then be transformed with high copy number plasmids for the expression of heterologous or homologous proteins.

20

Enzymes having characteristics which vary from available stock are required. In particular, enzymes having enhanced oxidation stability will be useful in extending the shelf life and bleach compatibility of proteases used in laundry products. Similarly, reduced oxidation stability would be useful in industrial processes that require the rapid and efficient quenching of enzymatic activity.

Modifying the pH-activity profiles of an enzyme would be useful in making the enzymes more efficient in a wide variety of processes, e.g. broadening the pH-activity profile of a protease would produce an enzyme more suitable for both alkaline and neutral laundry products. Narrowing the profile, particularly when combined with tailored substrate specificity, would make enzymes in a mixture more compatible, as will be further described herein.

35

Mutations of procaryotic carbonyl hydrolases (principally proteases but including lipases) will facilitate preparation of a variety of different hydrolases, particularly those having other modified properties such as  $K_m$ ,  $K_{cat}$ ,  $K_m/K_{cat}$  ratio and substrate specificity. These enzymes can then be tailored for the particular substrate which is anticipated to be present, for example in the preparation of peptides or for hydrolytic processes such as laundry uses.

Chemical modification of enzymes is known. For example, see I. Svendsen, 1976, "Carlsberg Res. Commun." 41 (5): 237-291. These methods, however, suffer from the disadvantages of being dependent upon the presence of convenient amino acid residues, are frequently nonspecific in that they modify all accessible residues with common side chains, and are not capable of reaching inaccessible amino acid residues without further processing, e.g. denaturation, that is generally not completely reversible in reinstituting activity. To the extent that such methods have the objective of replacing one amino acid residue side chain for another side chain or equivalent functionality, then mutagenesis promises to supplant such methods.

Predetermined, site-directed mutagenesis of tRNA synthetase in which a cys residue is converted to serine has been reported (G. Winter et al., 1982, "Nature" 299:756-758; A. Wilkinson et al., 1984, "Nature" 307:187-188). This method is not practical for large scale mutagenesis. It is an object herein to provide a convenient and rapid method for mutating DNA by saturation mutagenesis.

### 30 Summary

A method for producing procaryotic carbonyl hydrolase such as subtilisin and neutral protease in recombinant host cells is described in which expression vectors containing sequences which encode desired subtilisin or neutral protease, including the pro, 0992Y

pre, or prepro forms of these enzymes, are used to transform hosts, the host cultured and desired enzymes recovered. The coding sequence may correspond exactly to one found in nature, or may contain modifications which confer desirable properties on the protein that is produced, as is further described below.

The novel strains then are transformed with at least one DNA moiety encoding a polypeptide not otherwise expressed in the host strain, the transformed strains cultured and the polypeptide recovered from the culture. Ordinarily, the DNA moiety is a directed-mutant of a host *Bacillus* gene, although it may be DNA encoding a eucaryotic (yeast or mammalian) protein. The novel strains also serve as hosts for protein expressed from a bacterial gene derived from sources other than the host genome, or for vectors expressing these heterologous genes, or homologous genes from the host genome. In the latter event enzymes such as amylase are obtained free of neutral protease or subtilisin. In addition, it is now possible to obtain neutral protease in culture which is free of enzymatically active subtilisin, and vice-versa.

One may, by splicing the cloned genes for procaryotic carbonyl hydrolase into a high copy number plasmid, synthesize the enzymes in enhanced yield compared to the parental organisms. Also disclosed are modified forms of such hydrolases, including the pro and prepro zymogen forms of the enzymes, the pre forms, and directed mutations thereof.

A convenient method is provided for saturation mutagenesis, thereby enabling the rapid and efficient generation of a plurality of mutations at any one site within the coding region of a protein, comprising;

(a) obtaining a DNA moiety encoding at least a portion of said precursor protein;

0992Y

(b) identifying a region within the moiety;

(c) substituting nucleotides for those already existing within the region in order to create at least one restriction enzyme site unique to the moiety, whereby unique restriction sites 5' and 3' to the identified region are made available such that neither alters the amino acids coded for by the region as expressed;

(d) synthesizing a plurality of oligonucleotides, the 5' and 3' ends of which each contain sequences capable of annealing to the restriction enzyme sites introduced in step (c) and which, when ligated to the moiety, are expressed as substitutions, deletions and/or insertions of at least one amino acid in or into said precursor protein;

(e) digesting the moiety of step (c) with restriction enzymes capable of cleaving the unique sites; and

(f) ligating each of the oligonucleotides of step (d) into the digested moiety of step (e) whereby a plurality of mutant DNA moieties are obtained.

By the foregoing method or others known in the art, a mutation is introduced into isolated DNA encoding a procaryotic carbonyl hydrolase which, upon expression of the DNA, results in the substitution, deletion or insertion of at least one amino acid at a predetermined site in the hydrolase. This method is useful in creating mutants of wild type proteins (where the "precursor" protein is the wild type) or reverting mutants to the wild type (where the "precursor" is the mutant).

Mutant enzymes are recovered which exhibit oxidative stability and/or pH-activity profiles which differ from the precursor enzymes. Procaryotic carbonyl-hydrolases having varied  $K_m$ ,  $K_{cat}$ ,  $K_{cat}/K_m$  ratio and substrate specificity also are provided herein.



The mutant enzymes obtained by the methods herein are combined in known fashion with surfactants or detergents to produce novel compositions useful in the laundry or other cleaning arts.

5

Brief Description of the Drawings

Figure 1 shows the sequence of a functional B. amyloliquefaciens subtilisin gene.

10

In Figure 1A, the entire functional sequence for B. amyloliquefaciens, including the promoter and ribosome binding site, are present on a 1.5 kb fragment of the B. amyloliquefaciens genome.

15

Figure 1B shows the nucleotide sequence of the coding strand, correlated with the amino acid sequence of the protein. Promoter (p) ribosome binding site (rbs) and termination (term) regions of the DNA sequence are also shown.

20

Figure 2 shows the results of replica nitrocellulose filters of purified positive clones probed with Pool 1 (Panel A) and Pool 2 (Panel B) respectively.

25

Figure 3 shows the restriction analysis of the subtilisin expression plasmid (pS4). pBS42 vector sequences (4.5 kb) are shown in solid while the insert sequence (4.4 kb) is shown dashed.

30

Figure 5 shows the construction of the shuttle vector pBS42.

Figure 6 shows a restriction map for a sequence including the B. subtilis subtilisin gene.

35

0992Y

Figure 7 is the sequence of a functional B. subtilis subtilisin gene.

5 Figure 8 demonstrates a construction method for obtaining a deletion mutant of a B. subtilis subtilisin gene.

Figure 9 discloses the restriction map for a B. subtilis neutral protease gene.

10 Figure 10 is the nucleotide sequence for a B. subtilis neutral protease gene.

Figure 11 demonstrates the construction of a vector containing a B. subtilis neutral protease gene.

15 Figures 12, 13 and 16 disclose embodiments of the mutagenesis technique provided herein.

Figure 14 shows the enhanced oxidation stability of a subtilisin mutant.

Figure 15 demonstrates a change in the pH-activity profile of a subtilisin mutant when compared to the wild type enzyme.

25

#### Detailed Description

Procaryotic carbonyl hydrolases are enzymes which hydrolyze

30 compounds containing  $\begin{array}{c} \text{O} \\ \parallel \\ \text{C}-\text{X} \end{array}$  bonds in which X is oxygen or nitrogen. They principally include hydrolases, e.g. lipases and peptide hydrolases, e.g. subtilisins or metalloproteases. Peptide hydrolases include  $\alpha$ -aminoacylpeptide hydrolase, peptidylamino-acid hydrolase, acylamino hydrolase, serine carboxypeptidase, 35 metallocarboxypeptidase, thiol proteinase, carboxylproteinase and

metalloproteinase. Serine, metallo, thiol and acid proteases are included, as well as endo and exo-proteases.

5        Subtilisins are serine proteinases which generally act to cleave internal peptide bonds of proteins or peptides. Metalloproteases are exo- or endoproteases which require a metal ion cofactor for activity.

10       A number of naturally occurring mutants of subtilisin or neutral protease exist, and all may be employed with equal effect herein as sources for starting genetic material.

15       These enzymes and their genes may be obtained from many procaryotic organisms. Suitable examples include gram negative organisms such as E.coli or pseudomonas and gram positive bacteria such as micrococcus or bacillus.

20       The genes encoding the carbonyl hydrolase may be obtained in accord with the general method herein. As will be seen from the examples, this comprises synthesizing labelled probes having putative sequences encoding regions of the hydrolase of interest, preparing genomic libraries from organisms expressing the hydrolase, and screening the libraries for the gene of interest by hybridization to the probes. Positively hybridizing clones are then mapped and sequenced. The cloned genes are ligated into an expression vector (which also may be the cloning vector) with requisite regions for replication in the host, the plasmid transfected into a host for enzyme synthesis and the recombinant host cells cultured under conditions favoring enzyme synthesis, usually selection pressure such as is supplied by the presence of an antibiotic, the resistance to which is encoded by the vector. Culture under these conditions results in enzyme yields multifold greater than the wild type enzyme synthesis of the parent organism, even if it is the parent organism that is transformed.

35

"Expression vector" refers to a DNA construct containing a DNA sequence which is operably linked to a suitable control sequence capable of effecting the expression of said DNA in a suitable host. Such control sequences include a promoter to effect transcription, an optional operator sequence to control such transcription, a sequence encoding suitable mRNA ribosome binding sites, and sequences which control termination of transcription and translation. The vector may be a plasmid, a phage particle, or simply a potential genomic insert. Once transformed into a suitable host, the vector may replicate and function independently of the host genome, or may, in some instances, integrate into the genome itself. In the present specification, "plasmid" and "vector" are sometimes used interchangeably as the plasmid is the most commonly used form of vector at present. However, the invention is intended to include such other forms of expression vectors which serve equivalent functions and which are, or become, known in the art.

"Recombinant host cells" refers to cells which have been transformed or transfected with vectors constructed using recombinant DNA techniques. As relevant to the present invention, recombinant host cells are those which produce procaryotic carbonyl hydrolases in its various forms by virtue of having been transformed with expression vectors encoding these proteins. The recombinant host cells may or may not have produced a form of carbonyl hydrolase prior to transformation.

"Operably linked" when describing the relationship between two DNA regions simply means that they are functionally related to each other. For example, a presequence is operably linked to a peptide if it functions as a signal sequence, participating in the secretion of the mature form of the protein most probably involving cleavage of the signal sequence. A promoter is operably linked to a coding sequence if it controls the transcription of the sequence; a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation.

"Prohydrolase" refers to a hydrolase which contains additional N-terminal amino acid residues which render the enzyme inactive but, when removed, yield an enzyme. Many proteolytic enzymes are found in nature as translational proenzyme products and, in the absence of post-translational products, are expressed in this fashion.

"Presequence" refers to a signal sequence of amino acids bound to the N-terminal portion of the hydrolase which may participate in the secretion of the hydrolase. Presequences also may be modified in the same fashion as is described here, including the introduction of predetermined mutations. When bound to a hydrolase, the subject protein becomes a "prehydrolase". Accordingly, relevant prehydrolase for the purposes herein are presubtilisin and preprosubtilisin. Prehydrolases are produced by deleting the "pro" sequence (or at least that portion of the pro sequence that maintains the enzyme in its inactive state) from a prepro coding region, and then expressing the prehydrolase. In this way the organism excretes the active rather than proenzyme.

The cloned carbonyl hydrolase is used to transform a host cell in order to express the hydrolase. This will be of interest where the hydrolase has commercial use in its unmodified form, as for example subtilisin in laundry products as noted above. In the preferred embodiment the hydrolase gene is ligated into a high copy number plasmid. This plasmid replicates in hosts in the sense that it contains the well-known elements necessary for plasmid replication: a promoter operably linked to the gene in question (which may be supplied as the gene's own homologous promotor if it is recognized, i.e., transcribed, by the host), a transcription termination and polyadenylation region (necessary for stability of the mRNA transcribed by the host from the hydrolase gene) which is exogenous or is supplied by the endogenous terminator region of the hydrolase gene and, desirably, a selection gene such as an antibiotic resistance gene that enables continuous cultural maintenance of plasmid-infected host cells by growth in

0992Y

antibiotic-containing media. High copy number plasmids also contain an origin of replication for the host, thereby enabling large numbers of plasmids to be generated in the cytoplasm without chromosomal limitations. However, it is within the scope herein to  
 5 integrate multiple copies of the hydrolase gene into host genome. This is facilitated by bacterial strains which are particularly susceptible to homologous recombination. The resulting host cells are termed recombinant host cells.

10 Once the carbonyl hydrolase gene has been cloned, a number of modifications are undertaken to enhance the use of the gene beyond synthesis of the wild type or precursor enzyme. A precursor enzyme is the enzyme prior to its modification as described in this application. Usually the precursor is the enzyme as expressed by  
 15 the organism which donated the DNA modified in accord herewith. The term "precursor" is to be understood as not implying that the product enzyme was the result of manipulation of the precursor enzyme per se.

20 In the first of these modifications, the gene may be deleted from a recombination positive ( $rec^+$ ) organism containing a homologous gene. This is accomplished by recombination of an in vitro deletion mutation of the cloned gene with the genome of the organism. Many strains of organisms such as E.coli and Bacillus are  
 25 known to be capable of recombination. All that is needed is for regions of the residual DNA from the deletion mutant to recombine with homologous regions of the candidate host. The deletion may be within the coding region (leaving enzymatically inactive polypeptides) or include the entire coding region as long as  
 30 homologous flanking regions (such as promoters or termination regions) exist in the host. Acceptability of the host for recombination deletion mutants is simply determined by screening for the deletion of the transformed phenotype. This is most readily accomplished in the case of carbonyl hydrolase by assaying host  
 35 cultures for loss of the ability to cleave a chromogenic substrate otherwise hydrolyzed by the hydrolase.

Transformed hosts contained the protease deletion mutants are useful for synthesis of products which are incompatible with proteolytic enzymes. These hosts by definition are incapable of excreting the deleted proteases described herein, yet are

5 substantially normally sporulating. Also the other growth characteristics of the transformants are substantially like the parental organism. Such organisms are useful in that it is expected they will exhibit comparatively less inactivation of heterologous proteins than the parents, and these hosts do have growth

10 characteristics superior to known protease-deficient organisms. However, the deletion of neutral protease and subtilisin as described in this application does not remove all of the proteolytic activity of Bacillus. It is believed that intracellular proteases which are not ordinarily excreted extracellularly "leak" or diffuse

15 from the cells during late phases of the culture. These intracellular proteases may or may not be subtilisin or neutral protease as those enzymes are defined herein. Accordingly, the novel Bacillus strains herein are incapable of excreting the subtilisin and/or neutral protease enzymes which ordinarily are

20 excreted extracellularly in the parent strains. "Incapable" means not revertible to the wild type. Reversion is a finite probability that exists with the heretofore known protease-deficient, naturally occurring strains since there is no assurance that the phenotype of such strains is not a function of a readily revertible mutation,

25 e.g. a point mutation. This to be contrasted with the extremely large deletions provided herein.

The deletion mutant-transformed host cells herein are free of genes encoding enzymatically active neutral protease or subtilisin,

30 which genes are defined as those being substantially homologous with the genes set forth in Figs. 1, 7 or 10. "Homologous" genes contain coding regions capable of hybridizing under high stringency conditions with the genes shown in Figs. 1, 7 or 10.

The microbial strains containing carbonyl hydrolase deletion mutants are useful in two principal processes. In one embodiment they are advantageous in the fermentative production of products ordinarily expressed by a host that are desirably uncontaminated with the protein encoded by the deletion gene. An example is fermentative synthesis of amylase, where contaminant proteases interfere in many industrial uses for amylase. The novel strains herein relieve the art from part of the burden of purifying such products free of contaminating carbonyl hydrolases.

10

In a second principal embodiment, subtilisin and neutral protease deletion-mutant strains are useful in the synthesis of protein which is not otherwise encoded by the strain. These proteins will fall within one of two classes. The first class consists of proteins encoded by genes exhibiting no substantial pretransformation homology with those of the host. These may be proteins from other procaryotes but ordinarily are eucaryotic proteins from yeast or higher eucaryotic organisms, particularly mammals. The novel strains herein serve as useful hosts for expressible vectors containing genes encoding such proteins because the probability for proteolytic degradation of the expressed, non-homologous proteins is reduced.

The second group consists of mutant host genes exhibiting substantial pretransformation homology with those of the host. These include mutations of procaryotic carbonyl hydrolases such as subtilisin and neutral protease, as well as microbial (rennin, for example rennin from the genus Mucor). These mutants are selected in order to improve the characteristics of the precursor enzyme for industrial uses.

A novel method is provided to facilitate the construction and identification of such mutants. First, the gene encoding the hydrolase is obtained and sequenced in whole or in part. Then the sequence is scanned for a point at which it is desired to make a

0992Y



mutation (deletion, insertion or substitution) of one or more amino acids in the expressed enzyme. The sequences flanking this point are evaluated for the presence of restriction sites for replacing a short segment of the gene with an oligonucleotide pool which when  
 5 expressed will encode various mutants. Since unique restriction sites are generally not present at locations within a convenient distance from the selected point (from 10 to 15 nucleotides), such sites are generated by substituting nucleotides in the gene in such a fashion that neither the reading frame nor the amino acids encoded.  
 10 are changed in the final construction. The task of locating suitable flanking regions and evaluating the needed changes to arrive at two unique restriction site sequences is made routine by the redundancy of the genetic code, a restriction enzyme map of the gene and the large number of different restriction enzymes. Note  
 15 that if a fortuitous flanking unique restriction site is available, the above method need be used only in connection with the flanking region which does not contain a site.

Mutation of the gene in order to change its sequence to conform  
 20 to the desired sequence is accomplished by M13 primer extension in accord with generally known methods. Once the gene is cloned, it is digested with the unique restriction enzymes and a plurality of end termini-complementary oligonucleotide cassettes are ligated into the unique sites. The mutagenesis is enormously simplified by this  
 25 method because all of the oligonucleotides can be synthesized so as to have the same restriction sites, and no synthetic linkers are necessary to create the restriction sites.

The number of commercially available restriction enzymes having  
 30 sites not present in the gene of interest is generally large. A suitable DNA sequence computer search program simplifies the task of finding potential 5' and 3' unique flanking sites. A primary constraint is that any mutation introduced in creation of the restriction site must be silent to the final constructed amino acid  
 35 coding sequence. For a candidate restriction site 5' to the target  
 0992Y

codon a sequence must exist in the gene which contains at least all the nucleotides but for one in the recognition sequence 5' to the cut of the candidate enzyme. For example, the blunt cutting enzyme SmaI (CCC/GGG) would be a 5' candidate if a nearby 5' sequence  
5 contained NCC, CNC, or CCN. Furthermore, if N needed to be altered to C this alteration must leave the amino acid coding sequence intact. In cases where a permanent silent mutation is necessary to introduce a restriction site one may want to avoid the introduction of a rarely used codon. A similar situation for SmaI would apply  
10 for 3' flanking sites except the sequence NGG, GNG, or GGN must exist. The criteria for locating candidate enzymes is most relaxed for blunt cutting enzymes and most stringent for 4 base overhang enzymes. In general many candidate sites are available. For the codon-222 target described herein a BalI site (TGG/CCA) could have  
15 been engineered in one base pair 5' from the KpnI site. A 3' EcoRV site (GAT/ATC) could have been employed 11 base pairs 5' to the PstI site. A cassette having termini ranging from a blunt end up to a four base-overhang will function without difficulty. In retrospect, this hypothetical EcoRV site would have significantly shortened the  
20 oligonucleotide cassette employed (9 and 13 base pairs) thus allowing greater purity and lower pool bias problems. Flanking sites should obviously be chosen which cannot themselves ligate so that ligation of the oligonucleotide cassette can be assured in a single orientation.

25 The mutation per se need not be predetermined. For example, an oligonucleotide cassette or fragment is randomly mutagenized with nitrosoguanidine or other mutagen and then in turn ligated into the hydrolase gene at a predetermined location.

30 The mutant carbonyl hydrolases expressed upon transformation of the suitable hosts are screened for enzymes exhibiting desired characteristics, e.g. substrate specificity, oxidation stability, pH-activity profiles and the like.

35

A change in substrate specificity is defined as a difference between the  $K_{cat}/K_m$  ratio of the precursor enzyme and that of the mutant. The  $K_{cat}/K_m$  ratio is a measure of catalytic efficiency. Prokaryotic carbonyl hydrolases with increased or diminished  $K_{cat}/K_m$  ratios are described in the examples. Generally, the objective will be to secure a mutant having a greater (numerically larger)  $K_{cat}/K_m$  ratio for a given substrate, thereby enabling the use of the enzyme to more efficiently act on a target substrate. An increase in  $K_{cat}/K_m$  ratio for one substrate may be accompanied by a reduction in  $K_{cat}/K_m$  ratio for another substrate. This is a shift in substrate specificity, and mutants exhibiting such shifts have utility where the precursors are undesirable, e.g. to prevent undesired hydrolysis of a particular substrate in an admixture of substrates.

$K_{cat}$  and  $K_m$  are measured in accord with known procedures, or as described in Example 18.

Oxidation stability is a further objective which is accomplished by mutants described in the examples. The stability may be enhanced or diminished as is desired for various uses. Enhanced stability is effected by deleting one or more methionine, tryptophan, cysteine or lysine residues and, optionally, substituting another amino acid residue not one of methionine, tryptophan, cysteine or lysine. The opposite substitutions result in diminished oxidation stability. The substituted residue is preferably alanyl, but neutral residues also are suitable.

Mutants are provided which exhibit modified pH-activity profiles. A pH-activity profile is a plot of pH against enzyme activity and may be constructed as illustrated in Example 19 or by methods known in the art. It may be desired to obtain mutants with broader profiles, i.e., those having greater activity at certain pH than the precursor, but no significantly greater activity at any pH, or mutants with sharper profiles, i.e. those having enhanced

activity when compared to the precursor at a given pH, and lesser activity elsewhere.

The foregoing mutants preferably are made within the active site of the enzyme as these mutations are most likely to influence activity. However, mutants at other sites important for enzyme stability or conformation are useful. In the case of *Bacillus subtilisin* or its pre, prepro and pro forms, mutations at tyrosine-1, aspartate+32, asparagine+155, tyrosine+104, methionine+222, glycine+166, histidine+64, glycine+169, phenylalanine+189, serine+33, serine+221, tyrosine+217, glutamate+156 and/or alanine+152 produce mutants having changes in the characteristics described above or in the processing of the enzyme. Note that these amino acid position numbers are those assigned to *B. amyloliquefaciens* subtilisin as seen from Fig. 7. It should be understood that a deletion or insertion in the N-terminal direction from a given position will shift the relative amino acid positions so that a residue will not occupy its original or wild type numerical position. Also, allelic differences and the variation among various procaryotic species will result in positions shifts, so that position 169 in such subtilisins will not be occupied by glycine. In such cases the new positions for glycine will be considered equivalent to and embraced within the designation glycine+169. The new position for glycine+169 is readily identified by scanning the subtilisin in question for a region homologous to glycine+169 in Fig. 7.

One or more, ordinarily up to about 10, amino acid residues may be mutated. However, there is no limit to the number of mutations that are to be made aside from commercial practicality.

The enzymes herein may be obtained as salts. It is clear that the ionization state of a protein will be dependent on the pH of the surrounding medium, if it is in solution, or of the solution from which it is prepared, if it is in solid form. Acidic proteins are commonly prepared as, for example, the ammonium, sodium, or

potassium salts; basic proteins as the chlorides, sulfates, or phosphates. Accordingly, the present application includes both electrically neutral and salt forms of the designated carbonyl hydrolases, and the term carbonyl hydrolase refers to the organic structural backbone regardless of ionization state.

The mutants are particularly useful in the food processing and cleaning arts. The carbonyl hydrolases, including mutants, are produced by fermentation as described herein and recovered by suitable techniques. See for example K. Anstrup, 1974, Industrial Aspects of Biochemistry, ed. B. Spencer pp. 23-46. They are formulated with detergents or other surfactants in accord with methods known per se for use in industrial processes, especially laundry. In the latter case the enzymes are combined with detergents, builders, bleach and/or fluorescent whitening agents as is known in the art for proteolytic enzymes. Suitable detergents include linear alkyl benzene sulfonates, alkyl ethoxylated sulfate, sulfated linear alcohol or ethoxylated linear alcohol. The compositions may be formulated in granular or liquid form. See for example U.S. Patents 3,623,957; 4,404,128; 4,381,247; 4,404,115; 4,318,818; 4,261,868; 4,242,219; 4,142,999; 4,111,855; 4,011,169; 4,090,973; 3,985,686; 3,790,482; 3,749,671; 3,560,392; 3,558,498; and 3,557,002.

The following disclosure is intended to serve as a representation of embodiments herein, and should not be construed as limiting the scope of this application.

#### Glossary of Experimental Manipulations

In order to simplify the Examples certain frequently occurring methods will be referenced by shorthand phrases.

Plasmids are designated by a small p preceeded and/or followed by capital letters and/or numbers. The starting plasmids herein are commercially available, are available on an unrestricted basis, or can be constructed from such available plasmids in accord with published procedures.

"Klenow treatment" refers to the process of filling a recessed 3' end of double stranded DNA with deoxyribonucleotides complementary to the nucleotides making up the protruding 5' end of the DNA strand. This process is usually used to fill in a recessed end resulting from a restriction enzyme cleavage of DNA. This creates a blunt or flush end, as may be required for further ligations. Treatment with Klenow is accomplished by reacting (generally for 15 minutes at 15°C) the appropriate complementary deoxyribonucleotides with the DNA to be filled in under the catalytic activity (usually 10 units) of the Klenow fragment of E. coli DNA polymerase I ("Klenow"). Klenow and the other reagents needed are commercially available. The procedure has been published extensively. See for example T. Maniatis et al., 1982, Molecular Cloning, pp. 107-108.

"Digestion" of DNA refers to catalytic cleavage of the DNA with an enzyme that acts only at certain locations in the DNA. Such enzymes are called restriction enzymes, and the sites for which each is specific is called a restriction site. "Partial" digestion refers to incomplete digestion by a restriction enzyme, i.e., conditions are chosen that result in cleavage of some but not all of the sites for a given restriction endonuclease in a DNA substrate. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements as established by the enzyme suppliers were used. Restriction enzymes commonly are designated by abbreviations composed of a capital letter followed by other letters and then, generally, a number representing the microorganism from which each restriction enzyme originally was obtained. In general, about 1 µg

0992Y

of plasmid or DNA fragment is used with about 1 unit of enzyme in about 20  $\mu$ l of buffer solution. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37°C are  
5 ordinarily used, but may vary in accordance with the supplier's instructions. After incubation, protein is removed by extraction with phenol and chloroform, and the digested nucleic acid is recovered from the aqueous fraction by precipitation with ethanol. Digestion with a restriction enzyme infrequently is followed with  
10 bacterial alkaline phosphatase hydrolysis of the terminal 5' phosphates to prevent the two restriction cleaved ends of a DNA fragment from "circularizing" or forming a closed loop that would impede insertion of another DNA fragment at the restriction site. Unless otherwise stated, digestion of plasmids is not followed by 5'  
15 terminal dephosphorylation. Procedures and reagents for dephosphorylation are conventional (T. Maniatis et al., Id., pp. 133-134).

"Recovery" or "isolation" of a given fragment of DNA from a  
20 restriction digest means separation of the digest on 6 percent polyacrylamide gel electrophoresis, identification of the fragment of interest by molecular weight (using DNA fragments of known molecular weight as markers), removal of the gel section containing the desired fragment, and separation of the gel from DNA. This  
25 procedure is known generally. For example, see R. Lawn et al., 1981, "Nucleic Acids Res." 9:6103-6114, and D. Goeddel et al., (1980) "Nucleic Acids Res." 8:4057.

"Southern Analysis" is a method by which the presence of DNA  
30 sequences in a digest or DNA-containing composition is confirmed by hybridization to a known, labelled oligonucleotide or DNA fragment. For the purposes herein, Southern analysis shall mean separation of digests on 1 percent agarose and depurination as described by G. Wahl et al., 1979, "Proc. Nat. Acad. Sci. U.S.A." 76:3683-3687,  
35 transfer to nitrocellulose by the method of E. Southern, 1975, 0992Y

"J. Mol. Biol." 98:503-517, and hybridization as described by T. Maniatis et al., 1978, "Cell" 15:687-701.

5 "Transformation" means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or chromosomal integrant. Unless otherwise stated, the method used herein for transformation of E. coli is the  $\text{CaCl}_2$  method of Mandel et al., 1970, "J. Mol. Biol." 53:154, and for Bacillus, the method of Anagnostopoulous et al., 1961, "J. Bact." 81:791-746.

10

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments (T. Maniatis et al., Id., p. 146). Unless otherwise stated, ligation was accomplished using known buffers and conditions with 10 units of T4 DNA ligase ("ligase") per 0.5  $\mu\text{g}$  of approximately equimolar amounts of the DNA fragments to be ligated. Plasmids from the transformants were prepared, analyzed by restriction mapping and/or sequenced by the method of Messing, et al., 1981, "Nucleic Acids Res.", 9:309.

20

"Preparation" of DNA from transformants means isolating plasmid DNA from microbial culture. Unless otherwise stated, the alkaline/SDS method of Maniatis et al., Id. p. 90., was used.

25

"Oligonucleotides" are short length single or double stranded polydeoxynucleotides which were chemically synthesized by the method of Crea et al., 1980, "Nucleic Acids Res." 8:2331-2348 (except that mesitylene nitrotriazole was used as a condensing agent) and then purified on polyacrylamide gels.

30

All literature citations are expressly incorporated by reference.

35



## Example 1

Preparation of a Genomic DNA Library from *B. amyloliquefaciens*  
and Isolation of its Subtilisin Gene

5

The known amino acid sequence of the extracellular *B. amyloliquefaciens* permits the construction of a suitable probe mixture. The sequence of the mature subtilisin is included (along with the additional information contributed by the present work) in Figure 1. All codon ambiguity for the sequence of amino acids at position 117 through 121 is covered by a pool of eight oligonucleotides of the sequence AA( $\overset{C}{T}$ )AA( $\overset{C}{T}$ )ATGGA( $\overset{C}{T}$ )GT.

15

Chromosomal DNA isolated from *B. amyloliquefaciens* (ATCC No. 23844) as described by J. Marmur, "J. Mol. Biol.", 3:208, was partially digested by Sau 3A, and the fragments size selected and ligated into the BamH 1 site of dephosphorylated pBS42. (pBS42 is shuttle vector containing origins of replication effective both in *E. coli* and *Bacillus*. It is prepared as described in Example 4.) The Sau3A fragment containing vectors were transformed into *E. coli* K12 strain 294 (ATCC No. 31446) according to the method of M. Mandel, et al., 1970, "J. Mol. Bio." 53: 154 using 80-400 nanograms of library DNA per 250 $\mu$ L of competent cells.

20

Cells from the transformation mixture were plated at a density of  $1-5 \times 10^3$  transformants per 150mm plate containing LB medium + 12.5  $\mu$ g/ml chloramphenicol, and grown overnight at 37°C until visible colonies appeared. The plates were then replica plated onto BA85 nitrocellulose filters overlayed on LB/chloramphenicol plates. The replica plates were grown 10-12 hours at 37°C and the filters transferred to fresh plates containing LB and 150  $\mu$ g/ml spectinomycin to amplify the plasmid pool.

25

30

After overnight incubation at 37°C, filters were processed essentially as described by Grunstein and Hogness, 1975, "Proc. Natl. Acad. Sci. USA" 72: 3961-3965.

35

Natl. Acad. Sci. (USA)" 72: 3961. Out of approximately 20,000 successful transformants, 25 positive colonies were found. Eight of these positives were streaked to purify individual clones. 24 clones from each streak were grown in microtiter wells, stamped on to two replica filters, and probed as described above with either AACAA(<sup>C</sup>/<sub>T</sub>)ATGGA(<sup>C</sup>/<sub>T</sub>)GT(pool 1) or AATAA(<sup>C</sup>/<sub>T</sub>)ATGGA(<sup>C</sup>/<sub>T</sub>)GT(pool 2) which differ by only one nucleotide. As shown in Figure 2, pool 1 hybridized to a much greater extent to all positive clones than did pool 2, suggesting specific hybridization.

Four out of five miniplasmid preparations (Maniatis et al., Id.) from positive clones gave identical restriction digest patterns when digested with Sau3A or HincII. The plasmid isolated from one of these four identical colonies by the method of Maniatis et al., Id., had the entire correct gene sequence and was designated pS4. The characteristics of this plasmid as determined by restriction analysis are shown in Figure 3.

## Example 2

### Expression of the Subtilisin Gene

Bacillus subtilis I-168 (Catalog No. 1-A1, Bacillus Genetic Stock Center) was transformed with pS4 and a single chloramphenicol resistant transformant then grown in minimal medium. After 24 hours, the culture was centrifuged and both the supernatant (10-200  $\mu$ l) and pellet assayed for proteolytic activity by measuring the change in absorbance per minute at 412 nm using 1 ml of the chromogenic substrate succinyl-L-ala-ala-pro-phe-p-nitroanilide (0.2  $\mu$ M) in 0.1M sodium phosphate (pH 8.0) at 25°C. A B. subtilis I-168 culture transformed with pBS42 used as a control showed less than 1/200 of the activity shown by the pS4 transformed culture. Greater than 95 percent of the protease activity of the pS4 culture was present in the supernatant,

and was completely inhibited by treatment with phenylmethylsulfonyl fluoride (PMSF) but not by EDTA.

Aliquots of the supernatants were treated with PMSF and EDTA to inhibit all protease activity and analyzed by 12 percent SDS-PAGE according to the method of Laemmli, U.K., 1970 "Nature", 227: 680. To prepare the supernatants, 16  $\mu$ L of supernatant was treated with 1mM PMSF, 10 mM EDTA for 10 minutes, and boiled with 4  $\mu$ L of 5x concentrated SDS sample buffer minus  $\beta$ -mercaptoethanol. The results of Coomassie stain on runs using supernatants of cells transformed with pS4, pBS42, and untransformed B. amyloliquefaciens are shown in Figure 4. Lane 3 shows authentic subtilisin from B. amyloliquefaciens. Lane 2 which is the supernatant from pBS42 transformed B. subtilis, does not give the 31,000 MW band associated with subtilisin which is exhibited by Lane 1 from pS4 transformed hosts. The approximately 31,000 MW band result for subtilisin is characteristic of the slower mobility shown by the known M.W. 27,500 subtilisin preparations in general.

20

### Example 3

#### Sequencing of the B. amyloliquefaciens Subtilisin Gene

The entire sequence of an EcoRI-BamHI fragment (wherein the EcoRI site was constructed by conversion of the HincII site) of pS4 was determined by the method of F. Sanger, 1977, "Proc. Natl. Acad. Sci (USA)", 74:5463. Referring to the restriction map shown in Figure 3, the BamHI-PvuII fragment was found to hybridize with pool 1 oligonucleotides by Southern analysis. Data obtained from sequencing of this fragment directed the sequencing of the remaining fragments (e.g. PvuII-HincII and AvaI-AvaI). The results are shown in Figure 1.

Examination of the sequence confirms the presence of codons for the mature subtilisin corresponding to that secreted by the 0992Y

B. amyloliquefaciens. Immediately upstream from this sequence is a series of 107 codons beginning with the GTG start codon at -107. Codon -107 to approximately codon -75 encodes an amino acid sequence whose characteristics correspond to that of known signal sequences.

5 (Most such signal sequences are 18-30 amino acids in length, have hydrophobic cores, and terminate in a small hydrophobic amino acid.) Accordingly, examination of the sequence data would indicate that codons -107 to approximately -75 encode the signal sequence; the remaining intervening codons between -75 and -1 presumably encode a

10 prosequence.

#### Example 4

#### 15 Construction of pBS42

pBS42 is formed by three-way ligation of fragments derived from pUB110, pC194, and pBR322 (see Figure 5). The fragment from pUB110 is the approximately 2600 base pair fragment between the HpaII site at

20 1900 and the BamHI site at 4500 and contains an origin of replication operable in Bacillus: T. Grycztan, et al., 1978 "J. Bacteriol.", 134: 318 (1978); A. Jalanko, et al., 1981 "Gene", 14: 325. The BamHI site was tested with Klenow. The pBR322 portion is the 1100 base pair fragment between the PvuII site at 2067 and the Sau3A site at 3223

25 which contains the E. coli origin of replication: F. Bolivar, et al., 1977 "Gene", 2: 95; J. Sutcliffe, 1978, Cold Spring Harbor Symposium 43: I, 77. The pC194 fragment is the 1200 base pair fragment between the HpaII site at 973 and the Sau3A site at 2006 which contains the gene for chloramphenicol resistance expressible in both E. coli and B.

30 subtilis: S. Ehrlich, "Proc. Natl. Acad. Sci. (USA)", 74:1680; S. Horynuchi et al., 1982, "J. Bacteriol." 150: 815.

pBS42 thus contains origins of replication operable both in E. coli and in Bacillus and an expressible gene for chloramphenicol

35 resistance.

Example 5

Isolation and Sequencing of the *B. subtilis* Subtilisin Gene

5     *B. subtilis* I168 chromosomal DNA was digested with EcoRI and the  
 fragments resolved on gel electrophoresis. A single 6 kb fragment  
 hybridized to a [ $\alpha$ -<sup>32</sup>P] CTP nick translation - labelled fragment  
 obtained from the C-terminus of the subtilisin structural gene in pS4,  
 described above. The 6 kb fragment was electroluted and ligated into  
 10 pBS42 which had been digested with EcoRI and treated with bacterial  
 alkaline phosphatase. *E. coli* ATCC 31446 was transformed with the  
 ligation mixture and transformants selected by growth on LB agar  
 containing 12.5  $\mu$ g chloramphenicol/ml. Plasmid DNA was prepared from  
 a pooled suspension of 5,000 transformed colonies. This DNA was  
 15 transformed into *B. subtilis* BG84, a protease deficient strain, the  
 preparation of which is described in Example 8 below. Colonies which  
 produced protease were screened by plating on LB agar plus 1.5 percent  
 w/w Carnation powdered nonfat skim milk and 5  $\mu$ g chloramphenicol/ml  
 (hereafter termed skim milk selection plates) and observing for zones  
 20 of clearance evidencing proteolytic activity.

Plasmid DNA was prepared from protease producing colonies,  
 digested with EcoRI, and examined by Southern analysis for the  
 presence of the 6 kb EcoRI insert by hybridization to the  
 25 <sup>32</sup>P-labelled C-terminus fragment of the subtilisin structural gene  
 from *B. amyloliquefaciens*. A positive clone was identified and the  
 plasmid was designated pS168.1. *B. subtilis* BG84 transformed with  
 pS168.1 excreted serine protease at a level 5-fold over that produced  
 in *B. subtilis* I168. Addition of EDTA to the supernatants did not  
 30 affect the assay results, but the addition of PMSF  
 (phenylmethylsulfonyl fluoride) to the supernatants reduced protease  
 activity to levels undetectable in the assay described in Example 8  
 for strain BG84.

A restriction map of the 6.5 kb EcoRI insert is shown in Fig. 6. The subtilisin gene was localized to within the 2.5 kb KpnI-EcoRI fragment by subcloning various restriction enzyme digests and testing for expression of subtilisin in B. subtilis BG84. Southern analysis with the labelled fragment from the C-terminus of the B. amyloliquefaciens subtilisin gene as a probe localized the C-terminus of the B. subtilis gene to within or part of the 631 bp HincII fragment B in the center of this subclone (see Fig. 6). The tandem HincII fragments B, C, and D and HincII-EcoRI fragment E (Fig. 6) were ligated into the M13 vectors mp8 or mp9 and sequenced in known fashion (J. Messing et al., 1982, "Gene" 19:209-276) using dideoxy chain termination (F. Sanger et al., 1977, "Proc. Nat. Acad. Sci. U.S.A." 74:5463-5467). The sequence of this region is shown in Fig. 7. The first 23 amino acids are believed to be a signal peptide. The remaining 83 amino acids between the signal sequence and the mature coding sequence constitute the putative "pro" sequence. The overlined nucleotides at the 3' end of the gene are believed to be transcription terminator regions. Two possible Shine-Dalgarno sequences are underlined upstream from the mature start codon.

20

#### Example 6

#### 25 Manufacture of an Inactivating Mutation of the B. subtilis Subtilisin Gene

A two step ligation, shown in Fig. 8, was required to construct a plasmid carrying a defective gene which would integrate into the Bacillus chromosome. In the first step, pS168.1, which contained the 6.5 kb insert originally recovered from the B. subtilis genomic library as described in Example 5 above, was digested with EcoRI, the reaction products treated with Klenow, the DNA digested with HincII, and the 800 bp EcoRI-HincII fragment E (see Fig. 6) that contains, in part, the 5' end of the B. subtilis subtilisin gene, was recovered. This fragment was ligated into pJH101 (pJH101 is available from 0992Y

35

J. Hoch (Scripps) and is described by F.A. Ferrari et al., 1983, "J. Bact." 134:318-329) that had been digested with HincII and treated with bacterial alkaline phosphatase. The resultant plasmid, pIDV1, contained fragment E in the orientation shown in Fig. 8. In the second step, pS168.1 was digested with HincII and the 700 bp HincII fragment B, which contains the 3' end of the subtilisin gene, was recovered. pIDV1 was digested at its unique HincII site and fragment B ligated to the linearized plasmid, transformed in E. coli ATCC 31,446, and selected on LB plates containing 12.5 µg chloramphenicol/ml or 20 µg ampicillin/ml. One resulting plasmid, designated pIDV1.4, contained fragment B in the correct orientation with respect to fragment E. This plasmid pIDV1.4, shown in Fig. 8, is a deletion derivative of the subtilisin gene containing portions of the 5' and 3' flanking sequences as well.

B. subtilis BG77, a partial protease-deficient mutant ( $\text{Prt}^{+/-}$ ) prepared in Example 8 below was transformed with pIDV1.4. Two classes of chloramphenicol resistant ( $\text{Cm}^r$ ) transformants were obtained. Seventy-five percent showed the same level of proteases as BG77 ( $\text{Prt}^{+/-}$ ) and 25 percent were almost completely protease deficient ( $\text{Prt}^-$ ) as observed by relative zones of clearing on plates containing LB agar plus skim milk. The  $\text{Cm}^r \text{Prt}^-$  transformants could not be due to a single crossover integration of the plasmid at the homologous regions for fragment E or B because, in such a case, the gene would be uninterrupted and the phenotype would be  $\text{Prt}^{+/-}$ . In fact, when either of fragments E or B were ligated independently into pJH101 and subsequently transformed into B. subtilis BG77, the protease deficient phenotype was not observed. The  $\text{Cm}^r$  phenotype of  $\text{Cm}^r \text{Prt}^-$  pIDV1.4 transformants was unstable in that  $\text{Cm}^s \text{Prt}^-$  derivatives could be isolated from  $\text{Cm}^r \text{Prt}^-$  cultures at a frequency of about 0.1 percent after 10 generations of growth in minimal medium in the absence of antibiotic selection. One such derivative was obtained and designated BG2018. The deletion was transferred into IA84 (a BGSC strain carrying two auxotrophic mutations flanking the subtilisin gene) by PBS1 transduction. The derivative organism was designated BG2019.

Example 7

Preparation of a Genomic DNA Library from B. subtilis and  
Isolation of its Neutral Protease Gene

5

The partial amino acid sequence of a neutral protease of B. subtilis is disclosed by P. Levy et al. 1975, "Proc. Nat. Acad. Sci. USA" 72:4341-4345. A region of the enzyme (Asp Gln Met Ile Tyr Gly) was selected from this published sequence in which the least  
10 redundancy existed in the potential codons for the amino acids in the region. 24 combinations were necessary to cover all the potential coding sequences, as described below.

15           GA<sup>T</sup> CA<sup>G</sup> ATG AT<sup>T</sup> TA<sup>T</sup> C GG  
              C    A    A       C    C  
                          A  
          Asp   Gln   Met   Ile   Tyr   Gly

20 Four pools, each containing six alternatives, were prepared as described above in Example 1. The pools were labelled by phosphorylation with [ $\gamma$ -<sup>32</sup>P] ATP.

The labelled pool containing sequences conforming closest to a unique sequence in a B. subtilis genome was selected by digesting  
25 B. subtilis (1A72, Bacillus Genetic Stock Center) DNA with various restriction enzymes, separating the digests on an electrophoresis gel, and hybridizing each of the four probe pools to each of the blotted digests under increasingly stringent conditions until a single band was seen to hybridize. Increasingly stringent  
30 conditions are those which tend to disfavor hybridization, e.g., increases in formamide concentration, decreases in salt concentration and increases in temperature. At 37°C in a solution of 5x Denhardt's, 5x SSC, 50 mM NaPO<sub>4</sub> pH 6.8 and 20 percent formamide, only pool 4 would hybridize to a blotted digest. These  
35 were selected as the proper hybridization conditions to be used for the neutral protease gene and pool 4 was used as the probe.



A lambda library of B. subtilis strain BGSC 1-A72 was prepared in conventional fashion by partial digestion of the Bacillus genomic DNA by Sau3A, separation of the partial digest by molecular weight on an electrophoresis gel, elution of 15-20 kb fragments (R. Lawn et al., 1981, "Nucleic Acids Res." 9:6103-6114), and ligation of the fragments to BamHI digested charon 30 phage using a Packagene kit from Promega Biotec.

E. coli DP50supF was used as the host for the phage library, although any known host for Charon lambda phage is satisfactory. The E. coli host was plated with the library phage and cultured, after which plaques were assayed for the presence of the neutral protease gene by transfer to nitrocellulose and screening with probe pool 4 (Benton and Davis, 1977, "Science" 196:180-182). Positive plaques were purified through two rounds of single plaque purification, and two plaques were chosen for further study, designated λNPRG1 and λNPRG2. DNA was prepared from each phage by restriction enzyme hydrolysis and separation on electrophoresis gels. The separated fragments were blotted and hybridized to labelled pool 4 oligonucleotides. This disclosed that λNPRG1 contained a 2400 bp HindIII hybridizing fragment, but no 4300 EcoRI fragment, while λNPRG2 contained a 4300 bp EcoRI fragment, but no 2400 bp HindIII fragment.

The 2400 bp λNPRG1 fragment was subcloned into the HindIII site of pJH101 by the following method. λNPRG1 was digested by HindIII, the digest fractionated by electrophoresis and the 2400 bp fragment recovered from the gel. The fragment was ligated to alkaline phosphatase-treated HindIII digested pJH101 and the ligation mixture used to transform E. coli ATCC 31446 by the calcium chloride shock method of V. Hershfield et al., 1974, "Proc. Nat. Acad. Sci. (U.S.A.)" 79:3455-3459). Transformants were identified by selecting colonies capable of growth on plates containing LB medium plus 12.5 µg chloramphenicol/ml.

0992Y

Transformant colonies yielded several plasmids. The orientation of the 2400 bp fragment in each plasmid was determined by conventional restriction analysis (orientation is the sense reading or transcriptional direction of the gene fragment in relation to the reading direction of the expression vector into which it is ligated.) Two plasmids with opposite orientations were obtained and designated pNPRsubH6 and pNPRsubH1.

The 4300 bp EcoRI fragment of  $\lambda$ NPRG2 was subcloned into pBR325 by the method described above for the 2400 bp fragment except that  $\lambda$ NPRG2 was digested with EcoRI and the plasmid was alkaline phosphatase-treated, EcoRI-digested pBR325. pBR325 is described by F. Bolivar, 1978, "Gene" 4:121-136. Two plasmids were identified in which the 4300 bp insert was present in different orientations. These two plasmids were designated pNPRsubRI and pNPRsubRIb.

#### Example 8

##### 20 Characterization of B. subtilis Neutral Protease Gene

The pNPRsubH1 insert was sequentially digested with different restriction endonucleases and blot hybridized with labelled pool 4 in order to prepare a restriction map of the insert (for general procedures of restriction mapping see T. Maniatis *et al.*, Id., p. 377). A 430 bp RsaI fragment was the smallest fragment that hybridized to probe pool 4. The RsaI fragment was ligated into the SmaI site of M13 mp8 (J. Messing *et al.*, 1982, "Gene" 19:269-276 and J. Messing in *Methods in Enzymology*, 1983, R. Wu *et al.*, Eds., 101:20-78) and the sequence determined by the chain-terminating dideoxy method (F. Sanger *et al.*, 1977, "Proc. Nat. Acad. Sci. U.S.A." 74:5463-5467). Other restriction fragments from the pNPRsubH1 insert were ligated into appropriate sites in M13 mp8 or M13 mp9 vectors and the sequences determined. As required, dITP was used to reduce compression artifacts (D. Mills *et al.*, 1979, "Proc. 0992Y

Nat. Acad. Sci. (U.S.A.)" 76:2232-2235). The restriction map for the pNPRsubH1 fragment is shown in Fig. 9. The sequences of the various fragments from restriction enzyme digests were compared and an open reading frame spanning a codon sequence translatable into the amino and carboxyl termini of the protease (P. Levy et al., Id.) was determined. An open reading frame is a DNA sequence commencing at a known point which in reading frame (every three nucleotides) does not contain any internal termination codons. The open reading frame extended past the amino terminus to the end of the 2400 bp HindIII fragment. The 1300 bp BglII - HindIII fragment was prepared from pNPRsubR1b (which contained the 4300 bp EcoRI fragment of  $\lambda$ NPRG2) and cloned in M13 mp8. The sequence of this fragment, which contained the portion of the neutral protease leader region not encoded by the 2400 bp fragment of pNPRsubH1, was determined for 400 nucleotides upstream from the HindIII site.

The entire nucleotide sequence as determined for this neutral protease gene, including the putative secretory leader and prepro sequence, are shown in Fig. 10. The numbers above the line refer to amino acid positions. The underlined nucleotides in Fig. 10 are believed to constitute the ribosome binding (Shine-Dalgarno) site, while the overlined nucleotides constitute a potential hairpin structure presumed to be a terminator. The first 27 - 28 of the deduced amino acids are believed to be the signal for the neutral protease, with a cleavage point at ala-27 or ala-28. The "pro" sequence of a proenzyme structure extends to the amino-terminal amino acid (ala-222) of the mature, active enzyme.

A high copy plasmid carrying the entire neutral protease gene was constructed by (Fig. 11) ligating the BglII fragment of pNPRsubR1, which contains 1900 bp (Fig. 9), with the PvuII - HindIII fragment of pNPRsubH1, which contains 1400 bp. pBS42 (from Example 4) was digested with BamHI and treated with bacterial alkaline phosphatase to prevent plasmid recircularization. pNPRsubR1 was digested with BglII, the 1900 bp fragment was isolated

from gel electrophoresis and ligated to the open BamHI sites of pBS42. The ligated plasmid was used to transform E. coli ATCC 31446 by the calcium chloride shock method (V. Hershfield et al., Id.), and transformed cells selected by growth on plates containing LB medium with 12.5 µg/ml chloramphenicol. A plasmid having the Bgl II fragment in the orientation shown in Fig. 11 was isolated from the transformants and designated pNPRsubB1. pNPRsubB1 was digested (linearized) with EcoRI, repaired to flush ends by Klenow treatment and then digested with HindIII. The larger fragment from the HindIII digestion (containing the sequence coding for the amino terminal and upstream regions) was recovered.

The carboxyl terminal region of the gene was supplied by a fragment from pNPRsubH1, obtained by digestion of pNPRsubH1 with PvuII and HindIII and recovery of the 1400 bp fragment. The flush end PvuII and the HindIII site of the 1400 bp fragment was ligated, respectively, to the blunted EcoRI and the HindIII site of pNPRsubB1, as shown in Fig. 11. This construct was used to transform B. subtilis strain BG84 which otherwise excreted no proteolytic activity by the assays described below. Transformants were selected on plates containing LB medium plus 1.5 percent carnation powdered nonfat milk and 5µg/ml chloramphenicol. Plasmids from colonies that cleared a large halo were analyzed. Plasmid pNPR10, incorporating the structural gene and flanking regions of the neutral protease gene, was determined by restriction analysis to have the structure shown in Fig. 11.

B. subtilis strain BG84 was produced by N-methyl-N'-nitro-N-nitrosoguanidine (NTG) mutagenesis of B. subtilis I168 according to the general technique of Adelberg et al., 1965, "Biochem. Biophys. Res. Commun." 18:788-795. Mutagenized strain I168 was plated on skim milk plates (without antibiotic). Colonies producing a smaller halo were picked for further analysis. Each colony was characterized for protease production on skim milk plates and amylase production on starch plates. One such isolate, which was 0992Y

partially protease deficient, amylase positive and capable of sporulation, was designated BG77. The protease deficiency mutation was designated prt-77. The prt-77 allele was moved to a spo0A background by congression as described below to produce strain BG84,  
5 a sporulation deficient strain.

Table A

10	Strain	Relevant Genotype	origin
	I168	<u>trpC2</u>	
	JH703	<u>trpC2</u> , <u>pheA12</u> , <u>spo0AΔ677</u>	Trousdale <u>et al.</u> <sup>a</sup>
15	BG16	<u>purB6</u> , <u>metB5</u> , <u>leuA8</u> , <u>lys-21</u> , <u>hisA</u> , <u>thr-5</u> <u>sacA321</u>	Pb 1665
	BG77	<u>trpC2</u> , <u>prt-77</u>	NTG x I168
	BG81	<u>metB5</u> , <u>prt-77</u>	BG16 DNA x BG77
20	BG84	<u>spo0A677</u> , <u>prt-77</u>	JH703 DNA x BG81

<sup>a</sup> "Mol. Gen. Genetics" 173:61 (1979)

25 BG84 was completely devoid of protease activity on skim milk plates and does not produce detectable levels of either subtilisin or neutral protease when assayed by measuring the change in absorbance at 412 nm per minute upon incubation with 0.2 μg/ml succinyl (-L-ala-L-ala-L-pro-L-phe) p-nitroanilide (Vega) in 0.1 M  
30 sodium phosphate, pH 8, at 25°C. BG84 was deposited in the ATCC as deposit number 39382 on July 21, 1983. Samples for subtilisin assay were taken from late logarithmic growth phase supernatants of cultures grown in modified Schaeffer's medium (T. Leighton et al., 1971, "J. Biol. Chem." 246:3189-3195).

35

0992Y

Example 9

Expression of the Neutral Protease Gene.

5 BG84 transformed with pNPR10 was inoculated into minimal media supplemented with 0.1 percent casein hydrolysate and 10  $\mu$ g chloramphenicol and cultured for 16 hours. 0.1 ml of culture supernatant was removed and added to a suspension of 1.4 mg/ml Azocoll proteolytic substrate (Sigma) in 10 mM Tris-HCl, 100 mM NaCl, 10 pH 6.8 and incubating with agitation. Undigested substrate was removed by centrifugation and the optical density read at 505 nm. Background values of an Azocoll substrate suspension were subtracted. The amount of protease excreted by a standard protease-expressing strain, BG16 was used to establish an arbitrary 15 level of 100. The results with BG16, and with BG84 transformed with control and neutral protease gene-containing plasmids are shown in Table B in Example 12 below. Transformation of the excreted protease-devoid B. subtilis strain BG84 results in excretion of protease activity at considerably greater levels than in BG16, the 20 wild-type strain.

Example 10

25 Manufacture of an Inactivating Mutation of the Neutral Protease Gene

The two RsaI bounded regions in the 2400 bp insert of pNPRsubH1, totalling 527 bp, can be deleted in order to produce an incomplete 30 structural gene. The translational products of this gene are enzymatically inactive. A plasmid having this deletion was constructed as follows. pJH101 was cleaved by digestion with HindIII and treated with bacterial alkaline phosphatase. The fragments of the neutral protease gene to be incorporated into 35 linearized pJH101 were obtained by digesting pNPRsubH1 with HindIII

and RsaI, and recovering the 1200 bp HindIII-RsaI and 680 bp RsaI-HindIII fragments by gel electrophoresis. These fragments were ligated into linearized pJH101 and used to transform E. coli ATCC 31446. Transformants were selected on plates containing LB medium and 20 µg ampicillin/ml. Plasmids were recovered from the transformants and assayed by restriction enzyme analysis to identify a plasmid having the two fragments in the same orientation as in the pNPRsubH1 starting plasmid. The plasmid lacking the internal RsaI fragments was designated pNPRsubH1Δ.

10

#### Example 11

##### Replacement of the Neutral Protease Gene with a Deletion Mutant

15

Plasmid pNPRsubH1Δ was transformed into B. subtilis strain BG2019 (the subtilisin deleted mutant from Example 6) and chromosomal integrants were selected on skim milk plates. Two types of Cm<sup>r</sup> transformants were noted, those with parental levels of proteolysis surrounding the colony, and those with almost no zone of proteolysis. Those lacking a zone of proteolysis were picked, restreaked to purify individual colonies, and their protease deficient character on skim milk plates confirmed. One of the Cm<sup>r</sup>, proteolysis deficient colonies was chosen for further studies (designated BG2034). Spontaneous Cm<sup>s</sup> revertants of BG2034 were isolated by overnight growth in LB media containing no Cm, plating for individual colonies, and replica plating on media with and without Cm. Three Cm<sup>s</sup> revertants were isolated, two of which were protease proficient, one of which was protease deficient (designated BG2036). Hybridization analysis of BG2036 confirmed that the plasmid had been lost from this strain, probably by recombination, leaving only the deletion fragments of subtilisin and neutral protease.

35

0992Y

Example 12

Phenotype of Strains Lacking Functional Subtilisin and Neutral  
Protease

The growth, sporulation and expression of proteases was examined in strains lacking a functional gene for either the neutral or alkaline protease or both. The expression of proteases was examined by a zone of clearing surrounding a colony on a skim milk plate and by measurement of the protease levels in liquid culture supernatants (Table B). A strain (BG2035) carrying the subtilisin gene deletion, and showed a 30 percent reduction level of protease activity and a normal halo on milk plates. Strain BG2043, carrying the deleted neutral protease gene and active subtilisin gene, and constructed by transforming BG16 (Ex. 8) with DNA from BG2036 (Example 11), showed an 80 percent reduction in protease activity and only a small halo on the milk plate. Strain BG2054, considered equivalent to BG2036

Table B  
Effect of protease deletions on protease expression and sporulation.

	<u>Genotype<sup>a</sup></u>	<u>Protease activity<sup>b</sup></u>	<u>Percent Sporulation</u>
BG16	Wild type	100	40
BG2035	apr $\Delta$ 684	70	20
BG2043	nprE $\Delta$ 522	20	20
BG2054	apr $\Delta$ 684,nprE $\Delta$ 522	ND	45
BG84(pBS42)	spo0A $\Delta$ 677,prt-77	ND	--
BG84(pNPR10)	spo0A $\Delta$ 677,prt-77	3000	--

<sup>a</sup>Only the loci relevant to the protease phenotype are shown.

<sup>b</sup>Protease activity is expressed in arbitrary units, BG16 was assigned a level of 100. ND indicates the level of protease was not detectable in the assay used.



(Example 11) in that it carried the foregoing deletions in both genes, showed no detectable protease activity in this assay and no detectable halo on milk plates. The deletion of either or both of the protease genes had no apparent effect on either growth or sporulation. Strains carrying these deletions had normal growth rates on both minimal glucose and LB media. The strains sporulated at frequencies comparable to the parent strain BG16. Examination of morphology of these strains showed no apparent differences from strains without such deletions.

### Example 13

#### Site-specific Saturation Mutagenesis of the B. Amylolyticus Subtilisin Gene at Position 222; Preparation of the Gene for Cassette Insertion

pS4-5, a derivative of pS4 made according to Wells et al., "Nucleic Acids Res.", 1983, 11:7911-7924 was digested with EcoRI and BamHI, and the 1.5 kb EcoRI-BamHI fragment recovered. This fragment was ligated into replicative form M-13 mp9 which had been digested with EcoRI and BamHI (Sanger et al., 1980, "J. Mol. Biol." 143 161-178. Messing et al., 1981, "Nucleic Acids Research" 9, 304-321. Messing, J. and Vieira, J. (1982) Gene 19, 269-276). The M-13 mp9 phage ligations, designated M-13 mp9 SUBT, were used to transform E. coli strain JM101 and single stranded phage DNA was prepared from a two mL overnight culture. An oligonucleotide primer was synthesized having the sequence 5'-GTACAACGGTACCTCACGCAGCTGCAGGAGCGGCTGC-3'. This primer conforms to the sequence of the subtilis gene fragment encoding amino acids 216-232 except that the 10 bp of codons for amino acids 222-225 were deleted, and the codons for amino acids 220, 227 and 228 were mutated to introduce a KpnI site 5' to the met-222 codon and a PstI site 3' to the met+222 codon. See Fig. 12. Substituted nucleotides are denoted by asterisks, the underlined codons in line 2 represent the new restriction sites and the scored sequence in line 4

represents the inserted oligonucleotides. The primer (about 15  $\mu$ M) was labelled with [ $^{32}$ P] by incubation with [ $\gamma$ - $^{32}$ P]-ATP (10  $\mu$ L in 20  $\mu$ L reaction)(Amersham 5000 Ci/mmol, 10218) and  $T_4$  polynucleotide kinase (10 units) followed by non-radioactive ATP (100  $\mu$ M) to allow complete phosphorylation of the mutagenesis primer. The kinase was inactivated by heating the phosphorylation mixture at 68°C for 15 min.

The primer was hybridized to M-13 mp9 SUBT as modified from Norris *et al.*, 1983, "Nucleic Acids Res." 11, 5103-5112 by combining 5  $\mu$ L of the labelled mutagenesis primer (~3  $\mu$ M), ~1  $\mu$ g M-13 mp9 SUBT template, 1  $\mu$ L of 1  $\mu$ M M-13 sequencing primer (17-mer), and 2.5  $\mu$ L of buffer (0.3 M Tris pH 8, 40 mM  $MgCl_2$ , 12 mM EDTA, 10 mM DTT, 0.5 mg/ml BSA). The mixture was heated to 68°C for 10 minutes and cooled 10 minutes at room temperature. To the annealing mixture was added 3.6  $\mu$ L of 0.25 mM dGTP, dCTP, dATP, and dTTP, 1.25  $\mu$ L of 10 mM ATP, 1  $\mu$ L ligase (4 units) and 1  $\mu$ L Klenow (5 units). The primer extension and ligation reaction (total volume 25  $\mu$ L) proceeded 2 hours at 14°C. The Klenow and ligase were inactivated by heating to 68°C for 20 min. The heated reaction mixture was digested with BamHI and EcoRI and an aliquot of the digest was applied to a 6 percent polyacrylamide gel and radioactive fragments were visualized by autoradiography. This showed the [ $^{32}$ P] mutagenesis primer had indeed been incorporated into the EcoRI-BamHI fragment containing the now mutated subtilisin gene.

The remainder of the digested reaction mixture was diluted to 200  $\mu$ L with 10 mM Tris, pH 8, containing 1 mM EDTA, extracted once with a 1:1 (v:v) phenol/chloroform mixture, then once with chloroform, and the aqueous phase recovered. 15  $\mu$ L of 5M ammonium acetate (pH 8) was added along with two volumes of ethanol to precipitate the DNA from the aqueous phase. The DNA was pelleted by centrifugation for five minutes in a microfuge and the supernatant was discarded. 300  $\mu$ L of 70 percent ethanol was added to wash the DNA pellet, the wash was discarded and the pellet lyophilized.

pBS42 from example 4 above was digested with BamHI and EcoRI and purified on an acrylamide gel to recover the vector. 0.5 $\mu$ g of the digested vector, 50 $\mu$ M ATP and 6 units ligase were dissolved in 20  $\mu$ l of ligation buffer. The ligation went overnight at 14°C. The DNA  
5 was transformed into E. coli 294 rec<sup>+</sup> and the transformants grown in 4 ml of LB medium containing 12.5  $\mu$ g/ml chloramphenicol. Plasmid DNA was prepared from this culture and digested with KpnI, EcoRI and BamHI. Analysis of the restriction fragments showed 30-50 percent  
10 of the molecules contained the expected KpnI site programmed by the mutagenesis primer. It was hypothesized that the plasmid population not including the KpnI site resulted from M-13 replication before bacterial repair of the mutagenesis site, thus producing a heterogenous population of KpnI<sup>+</sup> and KpnI<sup>-</sup> plasmids in some of the transformants. In order to obtain a pure culture of the KpnI<sup>+</sup>  
15 plasmid, the DNA was transformed a second time into E. coli to clone plasmids containing the new KpnI site. DNA was prepared from 16 such transformants and six were found to contain the expected KpnI site.

20 Preparative amounts of DNA were made from one of these six transformants (designated p $\Delta$ 222) and restriction analysis confirmed the presence and location of the expected KpnI and PstI sites. 40  $\mu$ g of p $\Delta$ 222 were digested in 300  $\mu$ L of KpnI buffer plus 30  $\mu$ L KpnI (300 units) for 1.5 h at 37°C. The DNA was precipitated with  
25 ethanol, washed with 70 percent ethanol, and lyophilized. The DNA pellet was taken up in 200  $\mu$ L HindIII buffer and digested with 20  $\mu$ L (500 units) PstI for 1.5 h at 37°C. The aqueous phase was extracted with phenol/CHCl<sub>3</sub> and the DNA precipitated with ethanol. The DNA was dissolved in water and purified by polyacrylamide gel  
30 electrophoresis. Following electroelution of the vector band (120 v for 2 h at 0°C in 0.1 times TBE (Maniatis et al., Id.)) the DNA was purified by phenol/CHCl<sub>3</sub> extraction, ethanol precipitation and ethanol washing.

35

0992Y

Although pΔ222 could be digested to completion (>98 percent) by either KpnI or PstI separately, exhaustive double digestion was incomplete (<<50 percent). This may have resulted from the fact that these sites were so close (10 bp) that digestion by KpnI allowed "breathing" of the DNA in the vicinity of the PstI site, i.e., strand separation or fraying. Since PstI will only cleave double stranded DNA, strand separation could inhibit subsequent PstI digestion.

10

## Example 14

Ligation of Oligonucleotide Cassettes into the Subtilisin Gene

15     10 μM of four complementary oligonucleotide pools (A-D, Table 1 below) which were not 5' phosphorylated were annealed in 20 μl ligase buffer by heating for five minutes at 68°C and then cooling for fifteen minutes at room temperature. 1 μM of each annealed oligonucleotide pool, ~0.2 μg KpnI and PstI-digested pΔ222 obtained  
20     in Example 13, 0.5 mM ATP, ligase buffer and 6 units T<sub>4</sub> DNA ligase in 20 μL total volume was reacted overnight at 14°C to ligate the pooled cassettes in the vector. A large excess of cassettes (~300x over the pΔ222 ends) was used in the ligation to help prevent intramolecular KpnI-KpnI ligation. The reaction was diluted by  
25     adding 25 μL of 10 mM Tris pH 8 containing 1 mM EDTA. The mixture was reannealed to avoid possible cassette concatemer formation by heating to 68°C for five minutes and cooling for 15 minutes at room temperature. The ligation mixtures from each pool were transformed, separately into *E. coli* 294 rec<sup>+</sup> cells. A small aliquot from each  
30     transformation mixture was plated to determine the number of independent transformants. The large number of transformants indicated a high probability of multiple mutagenesis. The rest of the transformants (~200-400 transformants) were cultured in 4 ml of LB medium plus 12.5 μg chloramphenicol/ml. DNA was prepared from  
35     each transformation pool (A-D). This DNA was digested with KpnI,

0.1 µg was used to retransform E. coli rec<sup>+</sup> and the mixture was plated to isolate individual colonies from each pool. Ligation of the cassettes into the gene and bacterial repair upon transformation destroyed the KpnI and PstI sites. Thus, only pΔ222 was cut when the transformant DNA was digested with KpnI. The cut plasmid would not transform E. coli. Individual transformants were grown in culture and DNA was prepared from 24 to 26 transformants per pool for direct plasmid sequencing. A synthetic oligonucleotide primer having the sequence 5'-GAGCTTGATGTCATGGC-3' was used to prime the dideoxy sequencing reaction. The mutants which were obtained are described in Table C below.

Two codon+222 mutants (i.e., gln and ile) were not found after the screening described. To obtain these a single 25mer oligonucleotide was synthesized for each mutant corresponding to the top oligonucleotide strand in Figure 12. Each was phosphorylated and annealed to the bottom strand of its respective nonphosphorylated oligonucleotide pool (i.e., pool A for gln and pool D for ile). This was ligated into KpnI and PstI digested pΔ222 and processed as described for the original oligonucleotide pools. The frequency of appearance for single mutants obtained this way was 2/8 and 0/7 for gln and ile, respectively. To avoid this apparent bias the top strand was phosphorylated and annealed to its unphosphorylated complementary pool. The heterophosphorylated cassette was ligated into cut pΔ222 and processed as before. The frequency of appearance of gln and ile mutants was now 7/7 and 7/7, respectively.

The data in Table C demonstrate a bias in the frequency of mutants obtained from the pools. This probably resulted from unequal representation of oligonucleotides in the pool. This may have been caused by unequal coupling of the particular trimers over the mutagenesis codon in the pool. Such a bias problem could be remedied by appropriate adjustment of trimer levels during synthesis to reflect equal reaction. In any case, mutants which were not

isolated in the primary screen were obtained by synthesizing a single strand oligonucleotide representing the desired mutation, phosphorylating both ends, annealing to the pool of non-phosphorylated complementary strands and ligating into the cassette site. A biased heteroduplex repair observed for the completely unphosphorylated cassette may result from the fact that position 222 is closer to the 5' end of the upper strand than it is to the 5' end of the lower strand (see Figure 12). Because a gap exists at the unphosphorylated 5' ends and the mismatch bubble in the double stranded DNA is at position 222, excision repair of the top strand gap would more readily maintain a circularly hybridized duplex capable of replication. Consistent with this hypothesis is the fact that the top strand could be completely retained by selective 5' phosphorylation. In this case only the bottom strand contained a 5' gap which could promote excision repair. This method is useful in directing biased incorporation of synthetic oligonucleotide strands when employing mutagenic oligonucleotide cassettes.

20

## Example 15

Site-Specific Mutagenesis of the Subtilisin Gene at Position 166

The procedure of Examples 13-14 was followed in substantial detail, except that the mutagenesis primer differed (the 37 mer shown in Fig. 13 was used), the two restriction enzymes were SacI and XmaIII rather than PstI and KpnI and the resulting constructions differed, as shown in Fig. 13.

30

Bacillus strains excreting mutant subtilisins at position 166 were obtained as described below in Example 16. The mutant subtilisins exhibiting substitutions of ala, asp, gln, phe, his, lys, asn, arg, and val for the wild-type residue were recovered.

35

0992Y

## Example 16

Preparation of Mutant Subtilisin Enzymes

5     B. subtilis strain BG2036 obtained by the method of Example 11  
was transformed by the plasmids of Examples 14, 15 or 20 and by  
pS4-5 as a control. Transformants were plated or cultured in shaker  
flasks for 16 to 48 h at 37°C in LB media plus 12.5 µg/ml  
chloramphenicol. Mutant enzymatically active subtilisin was  
10 recovered by dialyzing cell broth against 0.01M sodium phosphate  
buffer, pH 6.2. The dialyzed broth was then titrated to pH 6.2 with  
1N HCl and loaded on a 2.5 x 2 cm column of CM cellulose (CM-52  
Whatman). After washing with 0.01M sodium phosphate, pH 6.2, the  
subtilisins (except mutants at position +222) were eluted with the  
15 same buffer made 0.08N in NaCl. The mutant subtilisins at position  
+222 were each eluted with 0.1M sodium phosphate, pH 7.0. The  
purified mutant and wild type enzymes were then used in studies of  
oxidation stability,  $K_m$ ,  $K_{cat}$ ,  $K_{cat}/K_m$  ratio, pH optimum, and  
changes in substrate specificity.

20

25

30

35

0992Y

Table C Oligonucleotide Pool Organization  
and Frequency of Mutants Obtained

	Pool	Amino Acids	Codon-222a	Frequencyb
5	A	asp met cys arg gln unexpected mutantsa	GAT ATG TGT AGA GAA	2/25 3/25 13/25 2/25 0/25 5/25
10	B	leu pro phe tyr his unexpected mutants	CTT CCT TTC TAC CAC	1/25 3/25 6/25 5/25 1/25 9/25
15	C	glu ala thr lys asn unexpected mutants	GAA GCT ACA AAA AAC	3/17 3/17 1/17 1/17 1/17 8/17
20	D	gly trp ile ser val unexpected mutants	GGC TGG ATC AGC GTT	1/23 8/23 0/23 1/23 4/23 9/23

a Codons were chosen based on frequent use in the cloned subtilisin gene sequence (Wells et al., 1983, Id.).

b Frequency was determined from single track analysis by direct plasmid sequencing.

c Unexpected mutants generally comprised double mutants with changes in codons next to 222 or at the points of ligation. These were believed to result from impurities in the obigonucleotide pools and/or erroneous repair of the gapped ends.



Example 17

Mutant Subtilisin Exhibiting Improved Oxidation Stability

5        Subtilisins having cysteine and alanine substituted at the 222 position for wild-type methionine (Example 16) were assayed for resistance to oxidation by incubating with various concentrations of sodium hypochloride (Clorox Bleach).

10        To a total volume of 400  $\mu$ l of 0.1M, pH 7,  $\text{NaPO}_4$  buffer containing the indicated bleach concentrations (Fig. 14) sufficient enzyme was added to give a final concentration of 0.016 mg/ml of enzyme. The solutions were incubated at 25°C for 10 min. and assayed for enzyme activity as follows: 120  $\mu$ l of either ala+222 or  
15        wild type, or 100  $\mu$ l of the cys+222 incubation mixture was combined with 890  $\mu$ l 0.1M tris buffer at pH 8.6 and 10  $\mu$ l of a sAAPFPN (Example 18) substrate solution (20 mg/ml in DMSO). The rate of increase in absorbance at 410 nm due to release of p-nitroaniline (Del Mar, E.G., et al., 1979 "Anal. Biochem." 99, 316-320) was  
20        monitored. The results are shown in Fig. 14. The alanine substitution produced considerably more stable enzyme than either the wild-type enzyme or a mutant in which a labile cysteine residue was substituted for methionine. Surprisingly, the alanine substitution did not substantially interfere with enzyme activity  
25        against the assay substrate, yet conferred relative oxidation stability on the enzyme. The serine+222 mutant also exhibited improved oxidation stability.

30        Example 18

Mutant Subtilisins Exhibiting Modified Kinetics and Substrate Specificity

35        Various mutants for glycine+166 were screened for modified

Kcat, Km and Kcat/Km ratios. Kinetic parameters were obtained by analysis of the progress curves of the reactions. The rate of reaction was measured as a function of substrate concentration. Data was analyzed by fitting to the Michaelis-Menton equation using the non-linear regression algorithm of Marquardt (Marquardt, D. W. 1963, "J. Soc. Ind. Appl. Math." 11, 431-41). All reactions were conducted at 25°C in 0.1M tris buffer, pH 8.6, containing benzoyl-L-Valyl-Glycyl-L-Arginyl-p-nitroanilide (BVGRpN; Vega Biochemicals) at initial concentrations of 0.0025 M to 0.00026 M (depending on the value of Km for the enzyme of interest - concentrations were adjusted in each measurement so as to exceed Km) or succinyl-L-Alanyl-L-Alanyl-L-Prolyl-L-Phenylalanyl-p-nitroanilide (sAAPFPN; Vega Biochemicals) at initial concentrations of 0.0010 M to 0.00028 M (varying as described for BVGRpN).

The results obtained in these experiments were as follows:

Table D

Substrate	Enzyme	Kcat (s <sup>-1</sup> )	Km (M)	Kcat/Km
sAAPFPN	gly-166(wild type)	37	1.4x10 <sup>-4</sup>	3 x 10 <sup>5</sup>
	ala+166	19	2.7x10 <sup>-5</sup>	7 x 10 <sup>5</sup>
	asp+166	3	5.8x10 <sup>-4</sup>	5 x 10 <sup>3</sup>
	glu+166	11	3.4x10 <sup>-4</sup>	3 x 10 <sup>4</sup>
	phe+166	3	1.4x10 <sup>-5</sup>	2 x 10 <sup>5</sup>
	hys+166	15	1.1x10 <sup>-4</sup>	1 x 10 <sup>5</sup>
	lys+166	15	3.4x10 <sup>-5</sup>	4 x 10 <sup>5</sup>
	asn+166	26	1.4x10 <sup>-4</sup>	2 x 10 <sup>5</sup>
	arg+166	19	6.2x10 <sup>-5</sup>	3 x 10 <sup>5</sup>
	val+166	1	1.4x10 <sup>-4</sup>	1 x 10 <sup>4</sup>
BVGRpN	Wild Type	2	1.1x10 <sup>-3</sup>	2 x 10 <sup>3</sup>
	asp+166	2	4.1x10 <sup>-5</sup>	5 x 10 <sup>4</sup>
	glu+166	2	2.7x10 <sup>-5</sup>	7 x 10 <sup>4</sup>
	asn+166	1	1.2x10 <sup>-4</sup>	8 x 10 <sup>3</sup>

The  $K_{cat}/K_m$  ratio for each of the mutants varied from that of the wild-type enzyme. As a measure of catalytic efficiency, these ratios demonstrate that enzymes having much higher activity against a given substrate can be readily designed and selected by screening in accordance with the invention herein. For example, A166 exhibits over 2 times the activity of the wild type on sAAPFpN.

This data also demonstrates changes in substrate specificity upon mutation of the wild type enzyme. For example, the  $K_{cat}/K_m$  ratio for the D166 and E166 mutants is higher than the wild type enzyme with the BVGpN substrate, but qualitatively opposite results were obtained upon incubation with sAAPFpN. Accordingly, the D166 and E166 mutants were relatively more specific for BVGRpN than for sAAPFpN.

#### Example 19

##### Mutant Subtilisin Exhibiting Modified pH-Activity Profile

The pH profile of the Cys+222 mutant obtained in Example 16 was compared to that of the wild type enzyme. 10  $\mu$ l of 60 mg/ml sAAPFpN in DMSO, 10  $\mu$ l of Cys+222 (0.18 mg/ml) or wild type (0.5 mg/ml) and 980  $\mu$ l of buffer (for measurements at pH 6.6, 7.0 and 7.6, 0.1M  $\text{NaPO}_4$  buffer; at pH 8.2, 8.6 and 9.2, 0.1M tris buffer; and at pH 9.6 and 10.0, 0.1M glycine buffer), after which the initial rate of change in absorbance at 410 nm per minute was measured at each pH and the data plotted in Fig. 15. The Cys+222 mutant exhibits a sharper pH optimum than the wild type enzyme.

#### Example 20

##### Site-Specific Mutagenesis of the Subtilisin Gene at Position 169

The procedure of Examples 13-14 was followed in substantial detail, except that the mutagenesis primer differed (the primer

0992Y

shown in Fig. 16 was used), the two restriction enzymes were KpnI and EcoRV rather than PstI and KpnI and the resulting constructions differed, as shown in Fig. 16.

5 Bacillus strains excreting mutant subtilisins at position 169 were obtained as described below in Example 16. The mutant subtilisins exhibiting substitutions of ala and ser for the wild-type residue were recovered and assayed for changes in kinetic features. The assay employed SAAPFpN at pH 8.6 in the same fashion  
10 as set forth in Example 18. The results were as follows:

Table E

Enzyme	Kcat (s <sup>-1</sup> )	Km (M)	Kcat/Km
15 ala+169	58	$7.5 \times 10^{-5}$	$8 \times 10^5$
ser+169	38	$8.5 \times 10^{-5}$	$4 \times 10^5$

# Example 21

## 20 Alterations in Specific Activity on a Protein Substrate

Position 166 mutants from Examples 15 and 16 were assayed for alteration of specific activity on a naturally occurring protein substrate. Because these mutant proteases could display altered  
25 specificity as well as altered specific activity, the substrate should contain sufficient different cleavage sites i.e., acidic, basic, neutral, and hydrophobic, so as not to bias the assay toward a protease with one type of specificity. The substrate should also contain no derivitized residues that result in the masking of  
30 certain cleavage sites. The widely used substrates such as hemoglobin, azocollogen, azocasein, dimethyl casein, etc., were rejected on this basis. Bovine casein,  $\alpha$  and  $\alpha_2$  chains, was chosen as a suitable substrate.

35

A 1 percent casein (w/v) solution was prepared in a 100 mM Tris buffer, pH 8.0, 10 mM EDTA. The assay protocol is as follows:

- 790  $\mu$ l 50 mM Tris pH 8.2
- 5        100  $\mu$ l 1 percent casein (Sigma) solution
- 10  $\mu$ l test enzyme (10-200  $\mu$ g).

This assay mixture was mixed and allowed to incubate at room temperature for 20 minutes. The reaction was terminated upon the addition of 100  $\mu$ l 100 percent trichloroacetic acid, followed by incubation for 15 minutes at room temperature. The precipitated protein was pelleted by centrifugation and the optical density of the supernatant was determined spectrophotometrically at 280 nm. The optical density is a reflection of the amount of unprecipitated, i.e., hydrolyzed, casein in the reaction mixture. The amount of casein hydrolysed by each mutant protease was compared to a series of standards containing various amounts of the wild type protease, and the activity is expressed as a percentage of the corresponding wild type activity. Enzyme activities were converted to specific activity by dividing the casein hydrolysis activity by the 280 nm absorbance of the enzyme solution used in the assay.

All of the mutants which were assayed showed less specific activity on casein than the wild type with the exception of Asn+166 which was 26 percent more active on casein than the wild type. The mutant showing the least specific activity was ile+166 at 0.184 of the wild type activity.

30

35

CLAIMS

1. A method of preparing a procaryotic carbonyl hydrolase which comprises: culturing a recombinant host cell  
5 transformed with an expression vector comprising the DNA sequence encoding the hydrolase, and recovering the hydrolase from the cell culture.
2. The method of claim 1 wherein the hydrolase is a  
10 protease, preferably a subtilisin or a metalloprotease, most preferably a subtilisin, the DNA sequence preferably encoding subtilisin in the form of prosubtilisin or preprosubtilisin.
- 15 3. The method of claim 1 wherein the recombinant host cell is a strain of Bacillus, preferably a Bacillus subtilis.
4. The method of claim 2 wherein the DNA sequence  
20 encoding subtilisin is operably linked to its native promoter, to a promoter homologous to the host which is other than the native promoter, or to a promoter which is heterologous to the host.
- 25 5. The process of claim 2 wherein the recombinant host cell was transformed with an effective expression vector for

the DNA sequence encoding the procaryotic protease operably linked to its signal sequence.

6. A composition comprising a procaryotic carbonyl  
5 hydrolase, preferably a *Bacillus* hydrolase, and a host microorganism transformed so as to be capable of expressing the hydrolase.

7. A composition comprising prepro-, pre- or procarbonyl  
10 hydrolase, preferably prosubtilisin, essentially free of cells which express said prepro-, pre- or procarbonyl hydrolase.

8. A liquid detergent composition comprising B.  
15 amyloliquefaciens subtilisin.

9. An expression vector for a procaryotic carbonyl  
hydrolase which comprises a DNA sequence encoding the hydrolase operably linked to a promoter compatible with a  
20 suitable host cell.

10. A recombinant expression vector comprising a DNA  
sequence encoding a prepro- or procarbonyl hydrolase, preferably subtilisin, operably linked to a promoter  
25 compatible with a suitable host cell; or a cell, preferably a strain of Bacillus, transformed by a said vector.

11. A method comprising:

(a) isolating a DNA moiety encoding a procaryotic carbonyl hydrolase;

5

(b) introducing a mutation into a predetermined region in the DNA which, upon expression of the DNA, results in the substitution, deletion or insertion of at least one amino acid at a predetermined site in the hydrolase;

10

and optionally

(c) transforming a suitable host with the mutated DNA of step (b) and recovering the expression product of the mutated DNA.

15

12. The method of claim 11 wherein the mutation is predetermined; preferably expressed as the substitution or insertion of a single amino acid.

20

13. The method of claim 11 wherein the DNA was isolated as a fragment of genomic DNA from an organism expressing the carbonyl hydrolase; the fragment preferably consisting essentially of the structural gene for the hydrolase.

25

14. The method of claim 11 wherein the hydrolase is a protein hydrolase or lipase, preferably *Bacillus subtilisin*,



prosubtilisin, preprosubtilisin, a metalloprotease or other peptide hydrolase.

15. The method of claim 14 wherein the hydrolase is
- 5 Bacillus subtilisin and the mutation is introduced into the subtilisin at aspartate+32, asparagine+155, tyrosine+104, methionine+222, glycine+166, histidine+64, serine+221, glycine+169, glutamate+156, serine+33, phenylalanine+189, tyrosine+217 and/or alanine+152.

10

16. The method of claim 14 wherein the hydrolase is prosubtilisin or preprosubtilisin and the mutation is introduced into the presubtilisin or preprosubtilisin at tyrosine-1.

15

17. The method of claim 11 wherein the mutation is expressed as a mutant carbonyl hydrolase exhibiting a change in one or more of the oxidation stability,  $K_m$ ,  $K_{cat}$ ,  $K_{cat}/K_m$  ratio, substrate specificity, specific activity or pH
- 20 optimum of the hydrolase.

18. The method of claim 14 wherein the hydrolase is a peptide hydrolase selected from  $\alpha$ -aminoacylpeptide hydrolase, peptidylamino-acid hydrolase, acylamino
- 25 hydrolase, serine carboxypeptidase, metallo-carboxypeptidase, thiol proteinase, carboxylproteinase or metalloproteinase.

19. DNA encoding a predetermined mutant of a procaryotic carbonyl hydrolase; or a vector capable of transforming a host cell to produce a mutant bacterial carbonyl hydrolase, which vector comprises such DNA and which, upon  
5 transformation of the host cell, results in expression of the mutant hydrolase; or a host cell transformed with a said vector.
20. A mutant carbonyl hydrolase of the kind obtained by  
10 the method of claim 11.
21. A composition comprising the hydrolase of claim 20 in combination with a detergent, preferably a liquid detergent or a detergent in granular form; optionally additionally  
15 comprising a builder, bleach or fluorescent whitening agent.
22. The composition of claim 11 wherein the detergent is a linear alkyl benzene sulfonate, alkyl ethoxylated sulfate, sulfated linear alcohol or ethoxylated linear alcohol.  
20
23. The method of claim 17 wherein a mutant exhibiting a different substrate specificity is recovered.
24. The method of claim 11 wherein the mutation is within  
25 the enzyme active site.

25. The method of claim 11 wherein the DNA is expressed in a bacterial host cell, preferably a Bacillus species.
26. The method of claim 11 wherein the mutation is  
5 expressed as the substitution of one or more methionine, tryptophan, cysteine or lysine residues by a substituent amino acid residue not one of methionine, tryptophan, cysteine or lysine, preferably alanine or serine.
- 10 27. The method of claim 17 wherein the mutation renders the mutant enzyme either less oxidation stable or more oxidation stable than the precursor enzyme.
28. DNA encoding a predetermined enzymatically active  
15 mutant of a precursor procaryotic carbonyl hydrolase, said mutant exhibiting a different substrate specificity, oxidation stability and/or pH-activity profile than the precursor enzyme; or a vector capable of transforming a host cell to produce a mutant enzyme, which vector comprises such  
20 DNA and which, upon transformation of the host cell, results in expression of the mutant enzyme; or a host cell transformed with such a vector.
29. A method comprising culturing a host cell of claim 28  
25 and recovering therefrom the mutant enzyme.

30. A substantially normally sporulating Bacillus which is incapable of excreting subtilisin or neutral protease.

5 31. The Bacillus of claim 30, preferably free of Bacillus strains capable of excreting subtilisin or neutral protease, and comprising a mutant neutral protease gene, preferably nonrevertible, which gene contains a deletion that results in no expression or, upon expression, an enzymatically inactive polypeptide.

10

32. The Bacillus of claim 30 which is (a) incapable of excreting neutral protease and (b) transformed with at least one DNA moiety encoding a polypeptide not otherwise expressed by the Bacillus, preferably a subtilisin mutant or  
15 a eucaryotic protein.

33. A vegetative-phase Bacillus culture which is essentially free of neutral protease, or which is essentially free of subtilisin.

20

34. A Bacillus culture free of any gene capable of expressing enzymatically active neutral protease, or free of any gene capable of excreting enzymatically active subtilisin.

25

35. A vector comprising a deletion mutated neutral protease gene or a deletion mutated subtilisin gene.

36. A method comprising culturing the Bacillus of claim 30 until a protein not subtilisin or neutral protease, preferably amylase, has accumulated in the culture, and recovering the protein.

5

37. A method comprising:

(a) obtaining a DNA moiety encoding at least a portion of said precursor protein;

10

(b) identifying a region within the moiety;

(c) substituting nucleotides for those already existing within the region in order to create at least one  
15 restriction enzyme site unique to the moiety, whereby unique restriction sites 5' and 3' to the identified region are made available such that neither alters the amino acids coded for by the region as expressed;

20 (d) synthesizing a plurality of oligonucleotides, the 5' and 3' ends of which each contain sequences capable of annealing to the restriction enzyme sites introduced in step (c) and which, when ligated to the moiety, are expressed as substitutions, deletions and/or insertions of at least one  
25 amino acid in or into said precursor protein;

(e) digesting the moiety of step (c) with restriction enzymes capable of cleaving the unique sites;

(f) ligating each of the oligonucleotides of step (d) into the digested moiety of step (e) whereby a plurality of mutant DNA moieties are obtained;

and optionally the further steps of

10 (g) expressing each of said moieties as a mutant protein in a suitable host;

(h) recovering the mutant proteins of step (g); and

15 (i) screening the step (h) mutant proteins for the desirable characteristic.

38. The method of claim 37 wherein the restriction enzyme sites are different.


20

39. The method of claim 37 wherein the oligonucleotides are less than about 50 bp.

25

Fig.1.

A.



B.

5' 5 3 4 5 3'

1 GGTCTACTAAATATTTATCCATACATACAAATTAATACACAGAAATCTGTCTATTGGTTATTCTGCAAAATGAAAAAAGGACAGGATAAAGA -107  
fMet GTG  
 P RBS PRE PRO MAT TERM

-100 -90  
 Arg Gly Lys Lys Val Trp Ile Ser Leu Leu Phe Ala Leu Ala Leu Ile Phe Thr Met Ala Phe Gly Ser Thr Ser  
 99 AGA GGC AAA AAA GTA TGG ATC AGT TTG CTG TTT GCT TTA GCG TTA ATC TTT ACG ATG GCG TTC GGC AGC ACA TCC

-30 -20 -10  
 Ser Ala Gln Ala Ala Gly Lys Ser Asn Gly Glu Lys Lys Tyr Ile Val Gly Phe Lys Gln Thr Met Ser Thr Met  
 174 TCT GCC CAG GCG GCA GGG AAA TCA AAC GGG GAA AAG AAA TAT ATT GTC GGG TTT AAA CAG ACA ATG AGC ACG ATG

-50 -40  
 Ser Ala Ala Lys Lys Lys Asn Val Ile Ser Glu Lys Gly Gly Lys Val Gln Lys Gln Phe Lys Tyr Val Asp Ala  
 249 AGC GGC GCT AAG AAG AAA GAT GTC ATT TCT GAA AAA GGC GGG AAA GTG CAA AAG CAA TTC AAA TAT GTA GAC GCA

-30 -20 -10  
 Ala Ser Ala Thr Leu Asn Glu Lys Ala Val Lys Glu Leu Lys Lys Asp Pro Ser Val Ala Tyr Val Glu Glu Asp  
 324 GCT TCA GCT ACA TTA AAC GAA AAA GCT GTA AAA GAA TTG AAA AAA GAC CCG AGC GTC GCT TAC GTT GAA GAA GAT

-10 1  
 His Val Ala His Ala Tyr Ala Gln Ser Val Pro Tyr Gly Val Ser Gln Ile Lys Ala Pro Ala Leu His Ser Gln  
 399 CAC GTA GCA CAT GCG TAC GCG CAG TCC GTG CCT TAC GGC GTA TCA CAA ATT AAA GCC CTT CCG CAT TCT CAA

20 30 40  
 Gly Tyr Thr Gly Ser Asn Val Lys Val Ala Val Ile Asn Ser Gly Ile Asp Ser Ser His Pro Asn Leu Lys Val  
 474 GGC TAC ACT GGA TCA AAT GTT AAA GTA GCG GTT ATC GAC AGC GGT ATC GAT TCT TCT CAT CCT GAT TTA AAG GTA

50 60 70  
 Ala Gly Gly Ala Ser Met Val Pro Ser Glu Thr Asn Pro Phe Gln Asp Asn Asn Ser His Gly Thr His Val Ala  
 549 GCA GGC GGA GGC AGC ATG GTT CCT TCT GAA ACA AAT CCT TTC CAA GAC AAC AAC TCT CAC GGA ACT CAC GTT GCC

70 80 90  
 Gly Thr Val Ala Ala Leu Asn Asn Ser Ile Gly Val Leu Gly Val Ala Pro Ser Ala Ser Leu Tyr Ala Val Lys  
 624 GGC ACA GTT GCG GCT CTT AAT AAC TCA ATC GGT GTA TTA GCG GTT GCG CCA AGC GCA TCA CTT TAC GCT GTA AAA

100 110 120  
 Val Leu Gly Ala Asp Gly Ser Gly Gln Tyr Ser Trp Ile Ile Asn Gly Ile Glu Trp Ala Ile Ala Asn Asn Met  
 699 GTT CTC GGT GCT GAC GGT TCC GGC CAA TAC AGC TGG ATC ATT AAC GGA ATC GAG TGG GCG ATC GCA AAC AAT ATG

120 130 140  
 Asp Val Ile Asn Met Ser Leu Gly Gly Pro Ser Gly Ser Ala Ala Leu Lys Ala Ala Val Asp Lys Ala Val Ala  
 774 GAC GTT ATT AAC ATG AGC CTC GGC GGA CCT TCT GGT TCT GCT GCT TTA AAA GCG GCA GTT GAT AAA GCC GTT GCA

150 160 170  
 Ser Gly Val Val Val Val Ala Ala Ala Gly Asn Glu Gly Thr Ser Gly Ser Ser Ser Thr Val Gly Tyr Pro Gly  
 849 TCC GGC GTC GTA GTG GTT GCG GCA GCC GGT AAC GAA GGC ACT TCC GGC AGC TCA AGC ACA GTG GGC TAC CCT GGT

170 180 190  
 Lys Tyr Pro Ser Val Ile Ala Val Gly Ala Val Asp Ser Ser Asn Gln Arg Ala Ser Phe Ser Ser Val Gly Pro  
 924 AAA TAC CCT TCT GTC ATT GCA GTA GGC GCT GTT GAC AGC AGC AAC CAA AGA GCA TCT TTC TCA AGC GTA GGA CCT

200 210 220  
 Glu Leu Asp Val Met Ala Pro Gly Val Ser Ile Gln Ser Thr Leu Pro Gly Asn Lys Tyr Gly Ala Tyr Asn Gly  
 999 GAG CTT GAT GTC ATG GCA CCT GGC GTA TCT ATC CAA AGC ACG CTT CCT GGA AAC AAA TAC GGG GCG TAC AAC GGT

220 230 240  
 Thr Ser Met Ala Ser Pro His Val Ala Gly Ala Ala Ala Leu Ile Leu Ser Lys His Pro Asn Trp Thr Asn Thr  
 1074 ACG TCA ATG GCA TCT CCG CAC GTT GCC GGA GCG GCT GCT TTG ATT CTT TCT AAG CAC CCG AAC TGG ACA AAC ACT

250 260 270  
 Gln Val Arg Ser Ser Leu Glu Asn Thr Thr Thr Lys Leu Gly Asp Ser Phe Tyr Tyr Gly Lys Gly Leu Ile Asn  
 1149 CAA GTC CRC AGC AGT TTA GAA AAC ACC ACT ACA AAA CTT GGT GAT TCT TTC TAC TAT GGA AAA GGG CTG ATC AAC


270 275  
 Val Gln Ala Ala Ala Gln AC  
 1274 GTA CAG GCG GCA GCT CAG TAA AACAATAAAACCGGCTTGCCCGCGCGGTTTTTATTATTTTCTCTCCGCGATGTTCAATCGCGTC

1316 ATAATCGACGGATGGCTCCCTCTGAAATTTTAACGAGAAACGGCGGTTGACCCGCTCARTCCGTAACGCGCAATCCTGAAACGCTCTCAATCGCGG

1416 CTTCGCGTTTCCGTCAGCTCAATGCCGTACGGTCGCGCGGTTTTCTGTATACCGGAGACGCGATTGTAATCGGATC

*Fig. 2.*

Panel A: probe 1  $[^{32}\text{P-AA}\text{C}\text{AA}\text{C}\text{ATGGA}\text{C}\text{GT}]$



Panel B: probe 2  $[^{32}\text{P-AA}\text{TAA}\text{C}\text{ATGGA}\text{C}\text{GT}]$






Fig.3.






Fig. 4.





6/16

Fig. 6.




Fig.7

1 GATATACCTAAATAGAGATAAAATCATCTCAAAAAAATGGGTCTACTAAAAATATTATTCATCTATTACAATAAATTCACAGAATAGTCTTTTAAGTAAG

101 TCTACTCTGAATTTTTTAAAGGAGAGGGTAAAGA GTG AGA AGC AAA AAA TTG TGG ATC AGC TTG TTG TTT GCG TTA ACG TTA

185 ATC TTT ACG ATG GCG TTC AGC AAC ATG TCT GCG CAG GCT GCC GGA AAA AGC AGT ACA GAA AAG AAA TAC ATT GTC

260 GGA TTT AAA CAG ACA ATG AGT GCC ATG AGT TCC GCC AAG AAA AAG GAT GTT ATT TCT GAA AAA GGC GGA AAG GTT

335 CAA AAG CAA TTT AAG TAT GTT AAC GCG GCC GCA GCA ACA TTG GAT GAA AAA GCT GTA AAA GAA TTG AAA AAA GAT

410 CCG AGC GTT GCA TAT GTG GAA GAA GAT CAT ATT GCA CAT GAA TAT GCG CAA TCT GTT CCT TAT GGC ATT TCT CAA

485 ATT AAA GCG CCG GCT CTT CAC TCT CAA GGC TAC ACA GGC TCT AAC GTA AAA GTA GCT GTT ATC GAC AGC GGA ATT

560 GAC TCT TCT CAT CCT GAC TTA AAC GTC AGA GGC GGA GCA AGC TTC GTA CCT TCT GAA ACA AAC CCA TAC CAG GAC

635 GGC AGT TCT CAC GGT ACG CAT GTA GCC GGT ACG ATT GCC GCT CTT AAT AAC TCA ATC GGT GTT CTG GGC GTT AGC

710 CCA AGC GCA TCA TTA TAT GCA GTA AAA GTG CTT GAT TCA ACA GGA AGC GGC CAA TAT AGC TGG ATT ATT AAC GGC

785 ATT GAG TGG GCC ATT TCC AAC AAT ATG GAT GTT ATC AAC ATG AGC CTT GGC GGA CCT ACT GGT TCT ACA GCG CTG

860 AAA ACA GTC GTT GAC AAA GCC GTT TCC AGC GGT ATC GTC GTT GCT GCC GCA GCC GGA AAC GAA GGT TCA TCC GGA

935 AGC ACA AGC ACA GTC GGC TAC CCT GCA AAA TAT CCT TCT ACT ATT GCA GTA GGT GCG GTA AAC AGC AGC AAC CAA

1010 AGA GCT TCA TTC TCC AGC GCA GGT TCT GAG CTT GAT GTG ATG GCT CCT GGC GTG TCC ATC CAA AGC ACA CTT CCT

1085 GGA GGC ACT TAC GGC GCT TAT AAC GGA ACG TCC ATG GCG ACT CCT CAC GTT GCC GGA GCA GCA GCG TTA ATT CTT

1160 TCT AAG CAC CCG ACT TGG ACA AAC GCG CAA GTC CGT GAT CGT TTA GAA AGC ACT GCA ACA TAT CTT GGA AAC TCT

1235 TTC TAC TAT GGA AAA GGG TTA ATC AAC GTA CAA GCA GCT GCA CAA TAA TAGTAAAAAGAGCAGGTTCTCTCCATACCTGCTTC

1318 TTTTATTTGTACGATCCTGATGTTCCGGCGCATCTCTTCTCTCCGATGTTGAATCCGTTCCATGATCGACGGATGGCTGCCTCTGAAATCTTC

1418 ACAAGCACCGGAGGATCAACCTGCTCAGCCCCGTACGGCCAAATCTGAAACGTTTAACTGACTGCTTCTGTTCTCTGTC





Fig.9.



CACATGAC TTGATCATC TTGATATAT TCACACAGG ACATATAT CAATTTCG TAGTATGT AGTTTGTG CAGTATCA TATATATG AATATAGT TTTCAGTAT  
 10 20 30  
 met gly leu gly lys leu ser val arg ala ala ser phe met ser leu pro gly val gln ala ala glu gly his gln leu lys  
 TTCAAAAGG GGAATTAT GTG GGT TTA GGT AAG AAA TTG TCT GTT GGT GTC GCT TCG TTT ATG AGT TTA TCA ATC AGC CTG CCA GGT GTT CAG GCT GCT GAA GGT CAT CAG CTT AAA  
 40 50 60 70  
 glu asn gln thr asn phe leu ser lys lys pro ile ala gln ser glu leu ser ala pro asn asp lys ala val lys gln phe leu lys lys asn ser asn ile phe lys gly asp pro  
 GAG AAT CAA ACA AAT TTC CTC TCC AAA AAG CCG ATT GGG CAA TCA GAA CTC TCT GCA CCA AAT GAC AGC GTC AAG CAG TTT TTG AAA AAG AAC AGC AAC ATT TTT AAA GGT GAC CCT  
 80 90 100 110  
 ser lys ser val lys leu val glu ser thr thr asp ala leu gly tyr lys his phe arg tyr ala pro val val asn gly val pro ile lys asp ser gln val ile val his val asp  
 TCC AAA AGC GTG AAG CTT GTT GAA AGC AGC ACT GAT GCC CTT GGA TAC AAG CAC TTT CGA TAT GCG CCT GTC GTT AAG GGA GTG CCA ATT AAA GAT TCG CAA GTG ATC GTT CAC GTC GAT  
 120 130 140 150  
 lys ser asp asn val tyr ala val asn gly leu leu his asn gln ser ala ala lys thr asp asn ser gln lys val ser ser glu lys ala leu ala leu ala phe lys ala ile gly  
 AAA TCC GAT AAT GTC TAT GCG GTC AAT GGT GAA TTA CAC AAT CAA TCT GCT GCA AAA ACA GAT AAC AGC CAA AAA GTC TCT GCA AAA CCG CTG GCA CTC GCT TTC AAA GGT ATC GGC  
 160 170 180 190  
 lys ser pro asp ala val ser asn gly ala ala lys asn ser asn lys ala glu leu lys ala ile glu thr lys asp gly ser tyr arg leu ala tyr asp val thr ile arg tyr val  
 AAA TCA CCA GAC GCT GTT TCT AAC GGA GCG GCC AAA AAC AGC AAT AAA GCC GAA TTA AAA GCG ATA GCA AAA GAC GGC AGC TAT COT CTT TAC GAC GTG AGT ATT CCG TAT GTC  
 200 210 220 230  
 glu pro glu pro ala asn trp glu val leu val asp ala glu thr gly ser ile leu lys gln gln asn lys val glu his ala ala thr gly ser gly thr thr leu lys gly ala  
 GAG CCT GAA CCG CCA GAC AAC TGG GAA GTC TTA GTT GAC GCC GAA ACA GGC AUC ATT TTA AAA CAG CAA AAT AAA GTA GAT GGC GGC ACT GGA AGC GGA ACA AGC CTA AAG GGC GCA  
 240 250 260 270  
 thr val pro leu asn ile ser tyr glu gly lys tyr val leu arg asp leu ser lys pro thr gly thr gln ile ile thr tyr asp leu gln asn arg gln ser arg leu pro gly  
 ACT GTT CTT TTG AAC ATC TCT TAT GAA GGC GGA AAA TAT GTT CTA AGA GAT CTT TCA AAA CCA ACA GGC ACC CAA ATC ACA TAT BAT TTG CAA AAC AGA CAA AGC GGC CTT CCG GGC  
 280 290 300 310  
 thr leu val ser ser thr thr lys thr phe thr ser ser gln arg ala ala val asp ala his tyr asn leu gly lys val tyr asp tyr phe tyr ser asn phe lys arg asn ser  
 ACG CTT GTC TCA AGC ACA AGC AAA ACA TTT ACA TCT TCA CAG CAG CCG GCA GCC GTT GAC GCA CAC TAT AAC CTC GGT AAA GTG TAC GAT TAT TTT TAT TCA AAC TTT AAA CAA AAC AGC  
 320 330 340 350  
 tyr asp asn lys gly ser lys ile val ser ser val his tyr gln thr gln tyr asn ala ala trp thr gly asp gln met ile tyr gly asp gly asp gly ser phe phe ser pro  
 TAT GAT AAC AAA GGC AGT AAA ATC GTT TCT TCC GTT CAC TAC GGC ACT CAA TAC AAT AAC GCT GCA TGG ACA GGA GAC CAG ATG ATT TAC GGT GAT GGC GAC GGT TCA TTC TTC TCT CCG  
 360 370 380 390  
 leu ser gly ser leu asp val thr ala his glu met thr his gly val thr gln glu thr ala asn leu ile tyr glu asn gln pro gly ala leu asn glu ser phe ser asp val phe  
 CTT TCC GGC TCA TTA GAT GTG ACA GCG CAT GAA ATG ACA CAT GGC GTC ACC CAA GAA ACA GCG AAC TTG ATT TAT GAA AAT CAG CCA GGT GCA TTA AAC GAG TCT TTC TCT GAC GTA TTC  
 400 410 420 430  
 gly tyr phe asn asp thr glu asp trp asp ile gly glu asp ile thr val ser gln pro ala leu arg ser leu ser leu ser pro thr lys tyr asn gln pro asp asn tyr ala asn tyr  
 GGG TAT TTT AAC GAT ACA GAA GAC TGG GAC ATC GGT GAA GAC ATT ACG GTC AGC CAG CCT GCT CTT CCG AGC CTG TCC AAC CCT ACA AAA TAC AAC CAG CCG CCT GAC AAT TAC GGC AAT TAC  
 440 450 460 470  
 arg asn leu pro asn thr asp glu gly asp tyr gly gly val his thr asn ser gly ile pro asn lys ala ala tyr asn gln thr lys leu gly val ser lys ser gln gln ile  
 CGA AAC CTT CCA AAC ACA GAT GAA GGC GAT TAT GGC GGT GTA CAC ACA AAC AGC GGA ATT CCA AAC ACC GCT TAC AAC ACC ATC ACA AAA CTT GGT GTA TCT AAA TCA CAG CAA ATC  
 480 490 500 510  
 tyr tyr arg ala leu thr thr pro ser thr phe lys asp ala lys ala leu ile gln ser ala arg asp leu tyr gly ser thr asp ala ala lys val glu ala  
 TAT TAC GGT GGA TTA ACA AGC TAC CTC AGC CCT TCT TCC ACG TTC AAA GAT GGC AAG GCA GCT CTC ATT CAG TCT GGC GGT GAC CTC TAC GGC TCA ACT GAT GGC GCT AAA GTT GAA GCA  
 520 530  
 ala trp asn ala val gly leu oc  
 GCC TGG AAT GCT GTT GGA TTG TAA TATTAGAAA AGCTGAGT CCTCAGGCT TTATTTGTA CATATCTG TTTCTCTC AGCTGAGG ACAGAGAT GCTGCTGTA GACAGAAAC GCTCTGAT  
 TTGCTAAG AGGATGAC CCGCACTGC GCATTATA AAGCTATG ATTGATCA CATATGAT AGAGATTC

Fig.10.




*Fig.11.*

Fig.12.

	codon:	220	222	230
Wild type amino acid sequence:		Ala	Tyr Asn Gly Thr Ser Met Ala Ser Pro His Val Ala Gly Ala Ala	
1. Wild type DNA sequence:		5'-GCG TAC AAC GGT ACG TCA ATG GCA TCT CCG CAC GTT GCC GGA GCG GCT-3'		
		3'-CGC ATG TTG CCA TGC AGT TAC CGT AGA GGC GTG CAA CGG CCT CGC CGA-5'		
2. Δp222 DNA sequence:		5'-GCG TAC AAC <u>GGT</u> ACC TCA-----CG CAC <u>GCT</u> <u>GCA</u> <u>GGA</u> GCG GCT-3'		
		3'-CGC ATG TTG CCA TGG AGT GC GTG CGA CGT CCT CGC CGA-5'		
			KpnI	PstI
3. Δp222 cut with KpnI and PstI:		5'-GCG TAC AAC GGT AC		* pGGA GCG GCT-3'
		3'-CGC ATG TTG Cp		A CGT CCT CGC CGA-5'
4. cut Δp222 ligated with oligo-nucleotide pools:			***	*
		5'-GCG TAC AAC GGT ACG TCA NNN GCA TCT CCG CAC GTT GCA GGA GCG GCT-3'		
		3'-CGC ATG TTG CCA TGC AGT NNN CGT AGA GGC GTG CAA CGT CCT CGC CGA-5'		

Fig.13.

Wild type amino acid sequence:	thr ser gly ser ser ser thr var gly tyr pro gly
1. Wild type DNA sequence:	5' ACT TCC GGC AGC TCA AGC ACA GTC GGC TAC CCT GGT 3' 3' TCA AGG CCG TCG AGT TCG TGT CAG CCG ATG GGA CCA 5'
2. Δp166	* 5' ACT TCC GGC AGC TCA A _ _ _ _ _ C CCG GGT 3' 3' TGA AGG CCC TCG AGT T _ _ _ _ _ G GGC CCA 5'
3. Δp166 cut with SacI and XmaIII	* 5' ACT TCC GGC AGC T 3' TGA AGG CCCp * 5' ACT TCC GGC AGC T 3' TGA AGG CCCp * 5' ACT TCC GGC AGC TCA AGC ACG GTC NNN TAT CCG GGT 3' 3' TGA AGG CCC TCG AGT TCG TGT CAG NNN ATA GGC CCA 5'
4. Cut Δp166 ligated with oligonucleotide pools	* 5' ACT TCC GGC AGC TCA AGC ACG GTC NNN TAT CCG GGT 3' 3' TGA AGG CCC TCG AGT TCG TGT CAG NNN ATA GGC CCA 5'
Mutagenesis primer 37mer	5' AA GGC ACT TCC GGC AGC TCA ACC CCG GTA AA TAC CCT 3'

OXIDATIVE STABILITY OF THE CYS-222  
AND ALA-222 MUTANTS

Fig. 14.





Fig. 15.



## G-169 Saturation Mutagenesis

**codon:**

**Wild type amino acid sequence:**

162

169

173

ser ser thr val gly tyr pro gly lys tyr pro ser

1. Wild type DNA sequence:

5'-TCA AGC ACA GTG GGC TAC CCT GGT AAA TAC CCT TCT-3'  
3'-AGT TCG TGT CAC CCG ATG GGA CCA TTT ATG GGA AGA-5'

2.  $\Delta p169$  DNA sequence:

5'-TCA AGC ACA GTG GGG TAC CCT-----GA TAT CCT TCT-3'  
3'-AGT TCG TGT CAC CCC ATG GGA-----CT ATA GGA AGA-5'

\*                  \*                  \*

**KpnI**                      **EcoRV**

3.  $\Delta p169$  cut with KpnI and EcoRV:

5'-TAC AGC ACA GTC GGG TAC	*	
3'-AGT TCG TGT CAC CCP		
		*
		pAT CCT TCT-3'
		TA GGA AGA-5'

4. cut  $\Delta$ p169 ligated with oligo-nucleotide pools:

5'-TAC AGC ACA GTG GGG TAC CCT NNN AAA TAT CCT TGT-5' \*

3'-AGT TCG TGT CAC CCC ATG GGA NNN TTT ATA GGA AGA-3' \*

**Mutagenesis primer for Δp169**

5'-AAG CAC AGT GGG GTA CCC TGA TAT CCT TCT GTC A-3'



European Patent  
Office

# EUROPEAN SEARCH REPORT

0130756

Application number

EP 84304252.4

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
P, X	EP - A2 - 0 092 182 (GENENTECH, INC.) * Claims 18,19 *	1	C 12 N 15/00 C 12 N 9/14 C 12 N 9/54
X	EP - A2 - 0 057 976 (CPC INTERNATIONAL INC.) * Claims 1,2,7,12 *	1,3	C 11 D 3/386 C 11 D 7/42// C 12 R 1/125
D, X	US - A - 3 790 482 (JONES et al.) * Claim *	8,21, 22	
D, X	US - A - 3 557 002 (MC CARTY) * Claims 1,5; column 3, lines 21-26 *	8,21	
D, A	NATURE, vol. 299, no. 5885, October 21, 1982, New York, London G. WINTER et al. "Redesigning enzyme structure by sitedirected mutagenesis: tyrosyl t RNA synthetase and ATP binding" pages 756-758 * Totality *	1,3,6, 11,12, 17,19, 20,23-26,28, 35	C 12 N C 11 D
D, A, P	NATURE, vol. 307, January 12, 1984, New York, London A.J. WILKINSON et al. "A large increase in enzyme-substrate affinity by protein engineering" pages 187-188 * Totality *	11,12, 17,20, 23,24	
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 17-09-1984	Examiner WOLF
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			